

An Optimization Approach to the Langberg-Médard Multiple Unicast Conjecture ^{*†}

Kai Cai

The University of Hong Kong
email: kcai@hku.hk

Guangyue Han

The University of Hong Kong
email: ghan@hku.hk

June 7, 2018

Abstract

The Langberg-Médard multiple unicast conjecture claims that for any strongly reachable k -pair network, there exists a multi-flow with rate $(1, 1, \dots, 1)$. In a previous work, through combining and concatenating the so-called elementary flows, we have constructed a multi-flow with rate at least $(\frac{8}{9}, \frac{8}{9}, \dots, \frac{8}{9})$ for any k . In this paper, we examine an optimization problem arising from this construction framework. We first show that our previous construction yields a sequence of asymptotically optimal solutions to the aforementioned optimization problem. And furthermore, based on this solution sequence, we propose a perturbation framework, which not only promises a better solution for any $k \bmod 4 \neq 2$ but also solves the optimization problem for the cases $k = 3, 4, \dots, 10$, accordingly yielding multi-flows with the largest rate to date.

1 Introduction

We consider a directed k -pair network $\mathcal{N} = (V, A, S, R)$, which consists of an underlying digraph $D = (V, A)$, k senders $S = \{s_1, s_2, \dots, s_k\} \subseteq V$ and k receivers $R = \{r_1, r_2, \dots, r_k\} \subseteq V$. Let $\overline{\mathcal{N}}$ denote the underlying undirected network of \mathcal{N} , where the orientation in \mathcal{N} is ignored. Throughout this paper, we assume that each arc in \mathcal{N} (and as a result, each edge in $\overline{\mathcal{N}}$) is of unit capacity. The network coding rate $\mathbf{R}_c(\mathcal{N})$ is a real vector (d_1, d_2, \dots, d_k) such that d_i is the transmission rate from s_i to r_i when using network coding along the orientation of \mathcal{N} , while the routing rate $\mathbf{R}_c(\overline{\mathcal{N}})$ is a real vector (d_1, d_2, \dots, d_k) such that there exists a feasible (s_1, s_2, \dots, s_k) - (r_1, r_2, \dots, r_k) multi-flow (see definition in Section 2.1) over \mathcal{N} .

One of the most fundamental problems in the theory of network coding is *the multiple unicast network coding conjecture* [8], or simply *the multiple unicast conjecture*, which states that for any $\overline{\mathcal{N}}$, the transmission rate achieved by any fractional network coding can be achieved by routing as well. Despite enhanced understanding in certain special cases, the

*A preliminary version [21] of this work has been presented in IEEE ISIT 2018.

†This research is partly supported by a grant from the Research Grants Council of the Hong Kong Special Administrative Region, China (Project No. 17301017).

conjecture has been doggedly resisting a series of attacks [10]-[15] and turned out to be one of the most hardest problems in network coding theory.

A weaker version of the conjecture, proposed by Langberg and Médard [15], focuses on a strongly reachable k -pair network \mathcal{N} and claims that (the “ \leq ” below should be interpreted in the pairwise sense)

$$\mathbf{R}_c(\mathcal{N}) \leq \mathbf{R}_r(\overline{\mathcal{N}}),$$

Here, a k -pair network \mathcal{N} is said to be *strongly reachable* if there exists an s_i - r_j directed path P_{s_i, r_j} for all feasible i, j , and the paths $P_{s_1, r_j}, P_{s_2, r_j}, \dots, P_{s_k, r_j}$ are *edge-disjoint* for each feasible j . Apparently, for a strongly reachable k -pair network \mathcal{N} , $\mathbf{R}_c(\mathcal{N}) \geq (1, 1, \dots, 1)$. So, if the multiple unicast conjecture is true, one will deduce that $\mathbf{R}_r(\overline{\mathcal{N}}) \geq (1, 1, \dots, 1)$. Langberg and Médard [15] showed that $\mathbf{R}_r(\overline{\mathcal{N}}) \geq (\frac{1}{3}, \frac{1}{3}, \dots, \frac{1}{3})$, which was further improved to $\mathbf{R}_r(\overline{\mathcal{N}}) \geq (\frac{8}{9}, \frac{8}{9}, \dots, \frac{8}{9})$ in [18] by way of combining and concatenating some so-called elementary flows.

In this paper, we will examine a sequence of optimization problems $\{\mathcal{P}_{S_k}\}$, whose optimal solutions will naturally give lower bounds on $\mathbf{R}_r(\overline{\mathcal{N}})$. We first prove that our construction in [18] yields $\{\mathcal{C}_k^*\}$, a sequence of asymptotically optimal solutions to $\{\mathcal{P}_{S_k}\}$. And furthermore, based on $\{\mathcal{C}_k^*\}$, we propose a perturbation framework to obtain $\{\mathcal{C}_k^{**}\}$, which promises a better solution than \mathcal{C}_k^* for any $k \bmod 4 \neq 2$ and further solves \mathcal{P}_{S_k} for $k = 3, 4, \dots, 10$, and thereby yielding multi-flows with the largest rate to date (see Section 6.4). Here we note that a prototypical version of the optimization problem \mathcal{P}_{S_k} was first proposed in [20]. The progress made in this work is due to a (rather) delicate study of structural and analytic aspects of the optimization problem, which include symmetries, asymptotics, behaviors upon perturbation and so on.

The rest of paper is organized as follows. In Section 2, we give some basic notions and facts in the theory of multi-flows, and we introduce the optimization problem \mathcal{P}_{S_k} and elaborate its connections with the theory of multi-flows. In Section 3, we investigate the symmetries of the optimization problem \mathcal{P}_{S_k} and give the limit of its optimal value as k tends to infinity. We introduce the so-called strong homogeneous flow \mathcal{C}_k^* in Section 4, where we first show that $\{\mathcal{C}_k^*\}$ is a sequence of asymptotically optimal solution for $\{\mathcal{P}_{S_k}\}$ and then prove that it gives the exact optimal solution when $k = 1, 2, 6, 10$. In Section 5, we propose a unified framework to perturb \mathcal{C}_k^* to obtain a better solution \mathcal{C}_k^{**} for any $k \bmod 4 \neq 2$. In Sections 6, we give \mathcal{C}_k^{**} for $k = 3, 4, 5, 7, 8, 9$ explicitly, and we further establish the optimality of these \mathcal{C}_k^{**} and their uniqueness for achieving optimality. Finally, the paper is concluded in Section 7.

2 Mathematical Preliminaries

2.1 Multi-Flow Basics

Let $D = (V, A)$ be a directed graph with vertex set V and arc set A . For an arc $a = (u, v) \in A$, let $tail(a)$, $head(a)$ denote its *tail* u , *head* v , respectively. For any $s, r \in V$, an s - r *flow* is a function $f : A \rightarrow \mathbb{R}$ satisfying the following *flow conservation law*: for any $v \notin \{s, r\}$,

$$excess_f(v) = 0, \tag{1}$$

where

$$\text{excess}_f(v) := \sum_{a \in A: \text{head}(a)=v} f(a) - \sum_{a \in A: \text{tail}(a)=v} f(a). \quad (2)$$

It is easy to see that $|\text{excess}_f(s)| = |\text{excess}_f(r)|$, which is called the *value (or rate)* of f . Note that the above definitions naturally give rise to a *fractional flow* on the underlying undirected graph of D , and it is not needed to differentiate an s - r flow from an r - s flow. This is different from Schrijver [16], where a flow must be a non-negative function.

There are two kinds of operations on the flows defined as above. Firstly, the set of all s - r flows naturally forms a linear space over \mathbb{R} ; particularly, for any two s - r flows f_1, f_2 and scalars $u, v \in \mathbb{R}$, and the function $f = uf_1 + vf_2$ is again an s - r flow. Secondly, let f be an s - t flow and g be a t - r flow such that

$$\text{excess}_f(t) = -\text{excess}_g(t).$$

Then by definition, $f+g$ is an s - r flow, which is called the *concatenation* of f and g . Adopting the notational convention in defining the concatenation of paths in [16], the concatenation of f and g will be denoted by fg .

An (s_1, s_2, \dots, s_k) - (r_1, r_2, \dots, r_k) *multi-flow* refers to a set of k flows $\mathcal{F} = \{f_i : i = 1, 2, \dots, k\}$, where each f_i is an s_i - r_i flow. We say \mathcal{F} has *rate* (d_1, d_2, \dots, d_k) , where $d_i := |\text{excess}_{f_i}(s_i)|$; and, for any given $a \in A$, we define $|\mathcal{F}|(a)$ as

$$|\mathcal{F}|(a) := \sum_{1 \leq i \leq k} |f_i(a)|. \quad (3)$$

The multi-flow $\mathcal{F} = \{f_i : i = 1, 2, \dots, k\}$ is said to be *feasible* with respect to capacity function c if $|\mathcal{F}|(a) \leq c(a)$ for all $a \in A$. Note that when $k = 1$, the multi-flow is just a flow f , and f is feasible if $|f(a)| \leq c(a)$ for all $a \in A$ (Here recall that we have assumed $c(a) \equiv 1$ in Section 1).

2.2 Elementary Flows

For a *strongly reachable* k -pair network $\mathcal{N} = (V, A, S, R)$, let $\mathbf{P} = \{P_{s_i, r_j}\}_{i,j=1}^k$ be a set of s_i - r_j directed paths, where the paths $P_{s_1, r_j}, P_{s_2, r_j}, \dots, P_{s_k, r_j}$ are *edge-disjoint* for each feasible j . For each $P_{s_i, r_j} \in \mathbf{P}$, define an s_i - r_j flow as follows:

$$f_{i,j}(a) = \begin{cases} 1, & a \in P_{s_i, r_j}, \\ 0, & \text{otherwise.} \end{cases}$$

Let $\mathbf{F} = \{f_{i,j} | 1 \leq i, j \leq k\}$, a set of *elementary flows* with respect to \mathbf{P} , which will be the “building blocks” for the multi-flow construction in this paper.

More specifically, let

$$\mathcal{C} = \left((c_{i,j}^{(1)}), (c_{i,j}^{(2)}), \dots, (c_{i,j}^{(k)}) \right)$$

be a k -tuple of $k \times k$ real matrices. And for $\ell = 1, 2, \dots, k$, consider $\mathcal{F} = \{f_1, f_2, \dots, f_k\}$, where

$$f_\ell = \sum_{i,j=1}^k c_{i,j}^{(\ell)} f_{i,j}. \quad (4)$$

The following theorem says that if \mathcal{C} satisfies certain conditions, then the constructed \mathcal{F} in (4) is also a multi-flow.

Theorem 2.1. $\mathcal{F} = \{f_1, f_2, \dots, f_k\}$ be an (s_1, s_2, \dots, s_k) - (r_1, r_2, \dots, r_k) multi-flow with rate $(1, 1, \dots, 1)$ if and only if each $(c_{i,j}^{(\ell)})$ satisfies:

$$\begin{aligned} 1) \quad & \sum_{j=1}^k c_{i,j}^{(\ell)} = 0, \text{ for all } i \neq \ell; \\ 2) \quad & \sum_{i=1}^k c_{i,j}^{(\ell)} = 0, \text{ for all } j \neq \ell; \\ 3) \quad & \sum_{i=1}^k \sum_{j=1}^k c_{i,j}^{(\ell)} \equiv 1. \end{aligned} \tag{5}$$

Proof. We only need to prove that f_ℓ is an s_ℓ - r_ℓ flow with rate 1 for any ℓ . To see this, Note that $excess_{f_\ell}(s_i) = \sum_{j=1}^k c_{i,j}^{(\ell)}$ and $excess_{f_\ell}(r_j) = \sum_{i=1}^k c_{i,j}^{(\ell)}$. Condition 1) implies that the conservation law is satisfied by all the senders except s_{i_0} ; Condition 2) implies that it is satisfied by all the receivers except r_{j_0} ; Condition 3) implies that the value of f_ℓ is 1. \square

2.3 The Optimization Problem $\mathcal{P}_{\mathfrak{S}_k}$

The optimization problem $\mathcal{P}_{\mathfrak{S}_k}$ to be introduced in this section is intimately connected with our multi-flow construction and will be the main subject of study in this paper.

Let

$$\mathfrak{S}_k := \left\{ \mathcal{C} = \left((c_{i,j}^{(1)}), (c_{i,j}^{(2)}), \dots, (c_{i,j}^{(k)}) \right) \mid \mathcal{C} \text{ satisfies (5)} \right\}.$$

Clearly, \mathfrak{S}_k is defined by a total of $2k - 1$ linearly independent constraints and is an affine subspace of \mathbb{R}^{k^3} with dimension $k(k - 1)^2$.

Throughout this paper, we will refer to a non-empty subset of $[k] \times [k]$ as a k -sample, where $[k] := \{1, 2, \dots, k\}$. For a given k -sample s and $\ell \in [k]$, we define a function $g_s^{(\ell)} : \mathfrak{S}_k \rightarrow \mathbb{R}$ as

$$g_s^{(\ell)}(\mathcal{C}) := \sum_{(i,j) \in s} c_{i,j}^{(\ell)},$$

based on which, we define $g_s : \mathfrak{S}_k \rightarrow \mathbb{R}$ as

$$g_s(\mathcal{C}) := \sum_{\ell=1}^k |g_s^{(\ell)}(\mathcal{C})|.$$

Furthermore, for a non-empty set \mathcal{S} of k -samples, we define

$$g_{\mathcal{S}}(\mathcal{C}) := \max_{s \in \mathcal{S}} \{g_s(\mathcal{C})\}.$$

Now, we are ready to introduce the optimization problem $\mathcal{P}_{\mathcal{S}}$ as follows:

$$\begin{aligned} & \text{minimize} && g_{\mathcal{S}}(\mathcal{C}) \\ & \text{subject to} && \mathcal{C} \in \mathfrak{S}_k. \end{aligned} \tag{\mathcal{P}_{\mathcal{S}}}$$

Note that g_S is continuous and lower bounded, and thereby its optimal value is achievable, i.e., there exists an *optimal solution (optimal point)* $\bar{\mathcal{C}} \in \mathfrak{S}_k$ such that

$$g_S(\bar{\mathcal{C}}) = \min_{\mathcal{C} \in \mathfrak{S}_k} g_S(\mathcal{C}).$$

The following theorem says that \mathcal{P}_S is a convex optimization problem. Though the theorem follows from a standard argument, we give its proof for the sake of completeness.

Lemma 2.2. *g_S is a convex function over \mathfrak{S}_k .*

Proof. Let $\mathcal{C}_1, \mathcal{C}_2 \in \mathfrak{S}_k$ and $\mathcal{C} = p\mathcal{C}_1 + q\mathcal{C}_2$, such that $p + q = 1$, $p, q \leq 0$. Clearly, $\mathcal{C} \in \mathfrak{S}_k$, and

$$\begin{aligned} g_S(\mathcal{C}) &= \max_{s \in S} \{g_s(\mathcal{C})\} \\ &= \max_{s \in S} \{|g_s^{(1)}(p\mathcal{C}_1 + q\mathcal{C}_2)| + \cdots + |g_s^{(k)}(p\mathcal{C}_1 + q\mathcal{C}_2)|\} \\ &\leq \max_{s \in S} \{p|g_s^{(1)}(\mathcal{C}_1)| + q|g_s^{(1)}(\mathcal{C}_2)| + \cdots + p|g_s^{(k)}(\mathcal{C}_1)| + q|g_s^{(k)}(\mathcal{C}_2)|\} \\ &= \max_{s \in S} \{p g_s(\mathcal{C}_1) + q g_s(\mathcal{C}_2)\} \\ &\leq p \max_{s \in S} \{g_s(\mathcal{C}_1)\} + q \max_{s \in S} \{g_s(\mathcal{C}_2)\} \\ &= p g_S(\mathcal{C}_1) + q g_S(\mathcal{C}_2), \end{aligned}$$

which completes the proof. \square

Definition 2.3. [**Strongly Reachable Sample Set**] The following set of k -samples, denoted by \mathcal{S}_k , is of particular interest for the consideration of strongly reachable k -pair networks:

$$\mathcal{S}_k := \{\{(i_1, j_1), \dots, (i_r, j_r)\} \subseteq [k] \times [k] \mid j_1 < j_2 < \cdots < j_r, 1 \leq r \leq k\}. \quad (6)$$

Put it differently, \mathcal{S}_k is composed of all the k -samples, each of which consists of elements whose two coordinates are distinct. Clearly, there are $(k+1)^k - 1$ samples in \mathcal{S}_k .

Example 2.4. It is easy to see that $\mathcal{S}_1 = \{\{(1, 1)\}\}$. And \mathcal{S}_2 is composed of 8 samples, $\{(1, 1)\}$, $\{(2, 1)\}$, $\{(1, 2)\}$, $\{(2, 2)\}$, $\{(1, 1), (1, 2)\}$, $\{(2, 1), (1, 2)\}$, $\{(1, 1), (2, 2)\}$, and $\{(2, 1), (2, 2)\}$. And one can verify that \mathcal{S}_3 contains 63 samples and \mathcal{S}_4 contains 624 samples.

Let $\mathcal{O}_{\mathcal{S}_k}$ denote the *optimal value* of $\mathcal{P}_{\mathcal{S}_k}$. The following theorem provide a key link connecting $\mathcal{P}_{\mathcal{S}_k}$ and $\mathcal{R}_r(\bar{\mathcal{N}})$, where \mathcal{N} is an arbitrary strongly reachable k -pair network.

Theorem 2.5. *For any strongly reachable k -pair network \mathcal{N} ,*

$$\mathcal{R}_r(\bar{\mathcal{N}}) \geq \left(\frac{1}{\mathcal{O}_{\mathcal{S}_k}}, \frac{1}{\mathcal{O}_{\mathcal{S}_k}}, \dots, \frac{1}{\mathcal{O}_{\mathcal{S}_k}} \right).$$

Proof. Let $\bar{\mathcal{C}} = \left((\bar{c}_{i,j}^{(1)}), (\bar{c}_{i,j}^{(2)}), \dots, (\bar{c}_{i,j}^{(k)}) \right)$ be an optimal point for $\mathcal{P}_{\mathcal{S}_k}$, that is to say, $\mathcal{O}_{\mathcal{S}_k} = g_{\mathcal{S}_k}(\bar{\mathcal{C}})$. And let $\mathcal{F} = \{f_1, f_2, \dots, f_k\}$ be the (s_1, s_2, \dots, s_k) - (r_1, r_2, \dots, r_k) multi-flow constructed from \mathbf{F} with coefficient matrices $\frac{1}{\mathcal{O}_{\mathcal{S}_k}} \mathcal{C}^* = \left(\frac{1}{\mathcal{O}_{\mathcal{S}_k}} (\bar{c}_{i,j}^{(1)}), \frac{1}{\mathcal{O}_{\mathcal{S}_k}} (\bar{c}_{i,j}^{(2)}), \dots, \frac{1}{\mathcal{O}_{\mathcal{S}_k}} (\bar{c}_{i,j}^{(k)}) \right)$. Clearly, \mathcal{F} achieves rate $\left(\frac{1}{\mathcal{O}_{\mathcal{S}_k}}, \frac{1}{\mathcal{O}_{\mathcal{S}_k}}, \dots, \frac{1}{\mathcal{O}_{\mathcal{S}_k}} \right)$.

To complete the proof, we only need to prove that \mathcal{F} is feasible. Towards this goal, for each arc a , let $\mathcal{P}(a) = \{P_{s_{i_1}, r_{j_1}}, P_{s_{i_1}, r_{j_1}}, \dots, P_{s_{i_\alpha(a)}, r_{j_\alpha(a)}}\} \subseteq \mathbf{P}$ be the set of all the paths passing through a . By the definition of a strongly reachable k -pair network, we have

$$s(a) := \{(i_1, j_1), (i_2, j_2), \dots, (i_\alpha(a), j_\alpha(a))\} \in \mathcal{S}_k.$$

Hence, we have

$$\begin{aligned} |\mathcal{F}|(a) &= |f_1(a)| + \dots + |f_k(a)| \\ &= \left| f_{s(a)}^{(1)} \left(\frac{1}{\mathcal{O}_{\mathcal{S}_k}} (\bar{c}_{i,j}^{(k)}) \right) \right| + \dots + \left| f_{s(a)}^{(k)} \left(\frac{1}{\mathcal{O}_{\mathcal{S}_k}} (\bar{c}_{i,j}^{(k)}) \right) \right| \\ &= f_{s(a)} \left(\frac{1}{\mathcal{O}_{\mathcal{S}_k}} \bar{c} \right) \\ &\leq 1, \end{aligned}$$

which implies that \mathcal{F} is feasible and thus completes the proof. \square

3 Symmetries and Asymptotics of $\mathcal{P}_{\mathcal{S}_k}$

Starting from this section, we will focus on solving the problem $\mathcal{P}_{\mathcal{S}_k}$. Apparently, the problem $\mathcal{P}_{\mathcal{S}_1}$ is trivial. In [20], we have shown that $\mathcal{P}_{\mathcal{S}_2}$ has optimal value 1, which is achieved by the unique optimal point

$$\left(\left(\begin{pmatrix} \frac{3}{4} & \frac{1}{4} \\ \frac{1}{4} & \frac{1}{4} \end{pmatrix}, \begin{pmatrix} \frac{-1}{4} & \frac{1}{4} \\ \frac{1}{4} & \frac{3}{4} \end{pmatrix} \right) \right).$$

However, the problem $\mathcal{P}_{\mathcal{S}_k}$, $k \geq 3$, becomes prohibitively complex and cannot be dealt with a case analysis as in [20]. Rather than a fixed $\mathcal{P}_{\mathcal{S}_k}$, this section is devoted to the asymptotics of $\{\mathcal{P}_{\mathcal{S}_k}\}$; more precisely we will establish $\lim_{k \rightarrow \infty} \mathcal{O}_{\mathcal{S}_k} = 9/8$, which, as will be shown in Section 4, can be achieved by a sequence of explicitly constructed solutions. As elaborated below, the key observation in deriving this result is some symmetric properties possessed by the optimal solutions of $\mathcal{P}_{\mathcal{S}_k}$.

3.1 Symmetries of $\mathcal{P}_{\mathcal{S}_k}$

In this section, we use $Sym(k)$ to denote the symmetric group on $[k]$. Note that a permutation in $Sym(k)$ can be written by a product of disjoint cyclic permutations (cycles), e.g., $\sigma = (15)(342) \in Sym(5)$. For any $\sigma \in Sym(k)$ and $\mathcal{C} = \left((c_{i,j}^{(1)}), (c_{i,j}^{(2)}), \dots, (c_{i,j}^{(k)}) \right) \in \mathfrak{S}_k$, we define

$$\sigma(\mathcal{C}) := \tilde{\mathcal{C}},$$

where $\tilde{\mathcal{C}} = \left((\tilde{c}_{i,j}^{(1)}), (\tilde{c}_{i,j}^{(2)}), \dots, (\tilde{c}_{i,j}^{(k)}) \right)$ with $\tilde{c}_{i,j}^{(\ell)} = c_{\sigma^{-1}(i), \sigma^{-1}(j)}^{(\sigma^{-1}(\ell))}$ for all feasible i, j, ℓ . Apparently, σ defines a one-to-one mapping from \mathfrak{S}_k to \mathfrak{S}_k .

Example 3.1. Let

$$\mathcal{C} = \left(\left(\begin{pmatrix} \frac{5}{9} & \frac{2}{9} & \frac{2}{9} \\ \frac{3}{9} & \frac{-1}{9} & \frac{-1}{9} \\ \frac{2}{9} & \frac{-1}{9} & \frac{-1}{9} \end{pmatrix}, \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} \frac{-1}{9} & \frac{-1}{9} & \frac{2}{9} \\ \frac{-1}{9} & \frac{-1}{9} & \frac{2}{9} \\ \frac{2}{9} & \frac{2}{9} & \frac{5}{9} \end{pmatrix} \right) \in \mathfrak{S}_3,$$

and let $\sigma_1 = (12)$ and $\sigma_2 = (123) \in \text{Sym}(3)$. Then, we have

$$\begin{aligned}\sigma_1(\mathcal{C}) &= \left(\left(\begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \begin{pmatrix} \frac{-1}{9} & \frac{2}{9} & \frac{-1}{9} \\ \frac{2}{9} & \frac{2}{9} & \frac{-1}{9} \\ \frac{-1}{9} & \frac{-1}{9} & \frac{2}{9} \end{pmatrix}, \begin{pmatrix} \frac{-1}{9} & \frac{-1}{9} & \frac{2}{9} \\ \frac{-1}{9} & \frac{-1}{9} & \frac{2}{9} \\ \frac{2}{9} & \frac{2}{9} & \frac{-1}{9} \end{pmatrix} \right), \\ \sigma_2(\mathcal{C}) &= \left(\left(\begin{pmatrix} \frac{5}{9} & \frac{2}{9} & \frac{2}{9} \\ \frac{2}{9} & \frac{-1}{9} & \frac{-1}{9} \\ \frac{2}{9} & \frac{-1}{9} & \frac{-1}{9} \end{pmatrix}, \begin{pmatrix} \frac{-1}{9} & \frac{2}{9} & \frac{-1}{9} \\ \frac{2}{9} & \frac{2}{9} & \frac{-1}{9} \\ \frac{-1}{9} & \frac{-1}{9} & \frac{2}{9} \end{pmatrix}, \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix} \right).\end{aligned}$$

Definition 3.2. [Fixed Point and Invariant Space] Let $\mathcal{C} \in \mathfrak{S}_k$. \mathcal{C} is called a *fixed point* if for all $\sigma \in \text{Sym}(k)$, $\sigma(\mathcal{C}) = \mathcal{C}$. The set of all the fixed points is called the *invariant space* of \mathfrak{S}_k , and will be denoted by $\mathfrak{S}_k^{\text{fix}}$.

The following theorem shows that $\mathfrak{S}_k^{\text{fix}}$ is in fact a 2-dimensional affine subspace of \mathfrak{S}_k .

Theorem 3.3. Let $\mathcal{C} = \left((c_{i,j}^{(1)}), (c_{i,j}^{(2)}), \dots, (c_{i,j}^{(k)}) \right) \in \mathfrak{S}_k$. Then, $\mathcal{C} \in \mathfrak{S}_k^{\text{fix}}$ if and only if \mathcal{C} takes the following form:

$$\mathcal{C} = \left(\left(\begin{pmatrix} x & a & a & \dots & a \\ a & y & b & \dots & b \\ a & b & y & \dots & b \\ \vdots & \vdots & & \ddots & \vdots \\ a & b & b & \dots & y \end{pmatrix}, \begin{pmatrix} y & a & b & \dots & b \\ a & x & a & \dots & a \\ b & a & y & \dots & b \\ \vdots & \vdots & & \ddots & \vdots \\ b & a & b & \dots & y \end{pmatrix}, \dots, \begin{pmatrix} y & b & b & \dots & a \\ b & y & b & \dots & a \\ b & b & y & \dots & a \\ \vdots & \vdots & & \ddots & \vdots \\ a & a & a & \dots & x \end{pmatrix} \right), \quad (7)$$

where $x + (k-1)a = 1$ and $y + a + (k-2)b = 0$.

Proof. Clearly, if \mathcal{C} takes the form in (7), then it is a fixed point. So we only need to prove the reverse direction.

Let \mathcal{J} denote the set of all the entries of \mathcal{C} , i.e., $\mathcal{J} := \{c_{i,j}^{(\ell)} : 1 \leq i, j, \ell \leq k\}$. Consider the group action of $\text{Sym}(k)$ on \mathcal{J} with $\sigma(c_{i,j}^{(\ell)}) = c_{\sigma(i),\sigma(j)}^{(\sigma(\ell))}$ for any $\sigma \in \text{Sym}(k)$ and any $c_{i,j}^{(\ell)} \in \mathcal{J}$. Clearly, under this group action, \mathcal{J} is partitioned into the following orbits: 1) $\mathcal{J}_1 := \{c_{i,j}^{(\ell)} : i = j = \ell\}$; 2) $\mathcal{J}_2 := \{c_{i,j}^{(\ell)} : i = j \neq \ell\}$; 3) $\mathcal{J}_3 := \{c_{i,j}^{(\ell)} : i = \ell \neq j\}$; 4) $\mathcal{J}_4 := \{c_{i,j}^{(\ell)} : j = \ell \neq i\}$; 5) $\mathcal{J}_5 := \{c_{i,j}^{(\ell)} : i \neq j, i \neq \ell, j \neq \ell\}$. It follows from the assumption that \mathcal{C} is a fixed point that $c_{i,j}^{(\ell)} = c_{\sigma(i),\sigma(j)}^{(\sigma(\ell))}$ for any feasible i, j, ℓ and any $\sigma \in \text{Sym}(k)$. In other words, the elements in a same orbit must have a same value, and therefore we can assume the existence of x, y, a_1, a_2, b such that

$$c_{i,j}^{(\ell)} = \begin{cases} x, & \text{if } c_{i,j}^{(\ell)} \in \mathcal{J}_1, \\ y, & \text{if } c_{i,j}^{(\ell)} \in \mathcal{J}_2, \\ a_1, & \text{if } c_{i,j}^{(\ell)} \in \mathcal{J}_3, \\ a_2, & \text{if } c_{i,j}^{(\ell)} \in \mathcal{J}_4, \\ b, & \text{if } c_{i,j}^{(\ell)} \in \mathcal{J}_5. \end{cases}$$

Note that from (5), we can deduce that for any ℓ , $\sum_{j=1}^k c_{\ell,j}^{(\ell)} = \sum_{i=1}^k c_{i,\ell}^{(\ell)}$, which implies $\sum_{j:j \neq \ell} c_{\ell,j}^{(\ell)} = \sum_{i:i \neq \ell} c_{i,\ell}^{(\ell)}$, or equivalently, $(k-1)a_1 = (k-1)a_2$. The proof of the theorem is then complete after writing a_1, a_2 as a . \square

For any $\sigma \in \text{Sym}(k)$ and any k -sample $s = \{(i_1, j_1), \dots, (i_r, j_r)\}$, we define

$$\sigma(s) := \{(\sigma(i_1), \sigma(j_1)), \dots, (\sigma(i_r), \sigma(j_r))\}.$$

It is easy to see that σ defines a one-to-one mapping from $2^{[k] \times [k]}$ to $2^{[k] \times [k]}$. For a quick example, let $s = \{(2, 1), (1, 2), (3, 3)\} \subseteq [3] \times [3]$ and let $\sigma_1 = (1, 3)$, $\sigma_2 = (2, 3)$. Then, $\sigma_1(s) = \{(2, 3), (3, 2), (1, 1)\}$, $\sigma_2(s) = \{(3, 1), (1, 3), (2, 2)\}$.

Together with Theorem 3.3, the following theorem drastically reduces the dimension of the parameter space for the purpose of solving $\mathcal{P}_{\mathcal{S}_k}$.

Theorem 3.4. $\mathcal{P}_{\mathcal{S}_k}$ has an optimal point within \mathfrak{S}_k^{fix} .

Proof. Suppose that $\bar{\mathcal{C}} \in \mathfrak{S}_k$ achieves the optimal value of $\mathcal{P}_{\mathcal{S}_k}$. By Definition 3.2, for any $\sigma \in \text{Sym}(k)$, $\sigma(\bar{\mathcal{C}})$ is an optimal point of $\mathcal{P}_{\mathcal{S}_k}$. Let

$$\hat{\mathcal{C}} = \frac{\sum_{\sigma \in \text{Sym}(k)} \sigma(\bar{\mathcal{C}})}{n!}$$

It is easy to see that for any $\sigma \in \text{Sym}(k)$, $\sigma(\hat{\mathcal{C}}) = \hat{\mathcal{C}}$. Hence, $\hat{\mathcal{C}} \in \mathfrak{S}_k^{fix}$. On the other hand, it follows from Lemma 2.2 that

$$g_{\mathcal{S}_k}(\hat{\mathcal{C}}) \leq \frac{\sum_{\sigma \in \text{Sym}(k)} g_{\mathcal{S}_k}(\sigma(\bar{\mathcal{C}}))}{n!}$$

and hence $\hat{\mathcal{C}}$ is an optimal point, which completes the proof. \square

3.2 Asymptotics of $\mathcal{P}_{\mathcal{S}_k}$

For a k -sample $s = \{(i_1, j_1), (i_2, j_2), \dots, (i_{\alpha(s)}, j_{\alpha(s)})\}$, we define the following multi-set:

$$\text{Ind}_s := \{i_1, j_1, i_2, j_2, \dots, i_{\alpha(s)}, j_{\alpha(s)}\},$$

where $\alpha(s)$ denotes the size of s . And for any $\ell = 1, 2, \dots, k$, denote by $m_{\text{Ind}_s}(\ell)$ the multiplicity of ℓ in Ind_s (if $\ell \notin \text{Ind}_s$, then $m_{\text{Ind}_s}(\ell) = 0$), and define

$$\beta(s) := |\{\ell : m_{\text{Ind}_s}(\ell) \neq 0\}|.$$

For a quick example, consider $s = \{(1, 1), (2, 2), (1, 3), (3, 4), (1, 6)\} \subseteq [6] \times [6]$. Then, $\text{Ind}_s = \{1, 1, 2, 2, 1, 3, 3, 4, 1, 6\}$, $m_{\text{Ind}_s}(1) = 4$, $m_{\text{Ind}_s}(2) = m_{\text{Ind}_s}(3) = 2$, $m_{\text{Ind}_s}(4) = m_{\text{Ind}_s}(6) = 1$, $m_{\text{Ind}_s}(5) = 0$ and $\alpha(s) = \beta(s) = 5$.

In this section, we characterize the asymptotics of $\{\mathcal{O}_k\}$ and thereby approximately “solve” $\mathcal{P}_{\mathcal{S}_k}$ for large k . We first recall the following theorem from [20].

Theorem 3.5. $\mathcal{O}_{\mathcal{S}_k} \leq \frac{9}{8}$ for $k \geq 3$.

By Theorem 3.4, there exists an optimal point $\mathcal{C}_k \in \mathfrak{S}_k^{fix}$ for $\mathcal{P}_{\mathcal{S}_k}$. Moreover, by Theorem 3.3, we can assume $\mathcal{C}_k = ((c_{i,j}^{(1)}), (c_{i,j}^{(2)}), \dots, (c_{i,j}^{(k)}))$ takes the form as in (7) with a, b, x, y replaced by a_k, b_k, x_k, y_k , respectively, to emphasize its dependence on k , that is,

$$c_{i,j}^{(\ell)} = \begin{cases} x_k, & \text{if } i = j = \ell, \\ y_k, & \text{if } i = j \neq \ell \\ a_k, & \text{if } i = \ell; j \neq \ell \text{ or } j = \ell; i \neq \ell, \\ b_k, & \text{if otherwise,} \end{cases} \quad (8)$$

where $x_k + (k-1)y_k = 1$ and $y_k + (k-2)b_k + a_k = 0$.

Lemma 3.6. For x_k, y_k, a_k, b_k defined in (8), we have

- 1) $y_k = O(\frac{1}{k^2})$;
- 2) $x_k = O(\frac{1}{k})$;
- 3) $a_k = \frac{1}{k} + O(\frac{1}{k^2})$;
- 4) $b_k = \frac{-1}{k^2} + O(\frac{1}{k^3})$.

Proof. By definition, for $s = \{(1, 1), (2, 2), \dots, (\ell, \ell)\} \in \mathcal{S}_k$ and the optimal point \mathcal{C}_k defined in (8), we have

$$g_s(\mathcal{C}_k) = \ell|x_k + (\ell - 1)y_k| + (k - \ell)|\ell y_k|. \quad (9)$$

Taking $\ell = k/2$ in (9) and applying Theorem 3.5, we have

$$\frac{k^2}{4}|y_k| \leq g_s(\mathcal{C}_k) \leq \mathcal{O}_{\mathcal{S}_k} = O(1),$$

which implies $y_k = O(\frac{1}{k^2})$. Hence 1) holds.

Taking $\ell = k$ in (9) and applying Theorem 3.5, we have

$$k|x_k + (k - 1)y_k| \leq \mathcal{O}_{\mathcal{S}_k} = O(1).$$

Then, from 1) we deduce that $y_k = O(\frac{1}{k^2})$, which further implies $x_k = O(\frac{1}{k})$ by the above equation. Hence 2) holds.

Noticing that $(k - 1)a_k = 1 - x_k$ and by 2), we have $a_k = \frac{1}{k} + O(\frac{1}{k^2})$. Hence 3) holds.

Noticing that $(k - 2)b_k = -a_k - y_k$ and by 3) and 1), we have $b_k = \frac{-1}{k^2} + O(\frac{1}{k^3})$. Hence 4) holds. \square

Now, we are ready to give the main result of this section.

Theorem 3.7.

$$\lim_{k \rightarrow \infty} \mathcal{O}_{\mathcal{S}_k} = \frac{9}{8}.$$

Proof. Let $s = \{(i_1, 1), (i_2, 2), \dots, (i_\ell, \ell)\} \in \mathcal{S}_k$ be such that $\{i_1, i_2, \dots, i_\ell\} = \{1, 2, \dots, \ell\}$ and $i_j \neq j$ for $j = 1, 2, \dots, \ell$. It can be easily verified that $Ind_s = \{1, 1, 2, 2, \dots, \ell, \ell\}$. Let \mathcal{C}_k be an optimal point of $\mathcal{P}_{\mathcal{S}_k}$ taking the form in (8). Then, by definition, we have

$$\begin{aligned} g_s(\mathcal{C}_k) &= \sum_{i=1}^k |m_{Ind_s}(i)a_k + (\ell - m_{Ind_s}(i))b_k| \\ &= \sum_{i \in Ind_s} |m_{Ind_s}(i)a_k + (\ell - m_{Ind_s}(i))b_k| + \sum_{i \notin Ind_s} |\ell b_k| \\ &= \sum_{i \in Ind_s} |2a_k + (\ell - 2)b_k| + \sum_{i \notin Ind_s} |\ell b_k| \\ &= \ell|2a_k + (\ell - 2)b_k| + (k - \ell)|\ell b_k|. \end{aligned}$$

It then follows from Lemma 3.6 that $2a_k + (\ell - 2)b_k > 0$ and $b_k < 0$, and furthermore,

$$\begin{aligned}
g_s(\mathcal{C}_k) &= \ell|2a_k + (\ell - 2)b_k| + (k - \ell)|\ell b_k| \\
&= \ell(2a_k + (2\ell - k - 2)b_k) \\
&= \ell \left(\frac{2}{k} + O\left(\frac{1}{k^2}\right) - \frac{2\ell}{k^2} + \frac{1}{k} + O\left(\frac{1}{k^2}\right) \right) \\
&= \frac{\ell}{k^2}(3k - 2\ell + O(1)).
\end{aligned} \tag{10}$$

Now, setting $\ell = \frac{3k}{4} + O(1)$ in Equation (10), we have

$$g_s(\mathcal{C}_k) = \frac{1}{k^2} \left(\frac{3k}{4} + O(1) \right) \left(\frac{3k}{2} + O(1) \right) = \frac{9}{8} + O\left(\frac{1}{k}\right).$$

Hence, $\mathcal{O}_{S_k} \geq g_s(\mathcal{C}_k) = \frac{9}{8} + O\left(\frac{1}{k}\right)$. On the other hand, by Lemma 3.5, we have that $\mathcal{O}_{S_k} \leq \frac{9}{8}$, which immediately implies that

$$\lim_{k \rightarrow \infty} \mathcal{O}_{S_k} = \frac{9}{8}.$$

□

4 The Strong Homogeneous Flow \mathcal{C}_k^*

We introduce in this section a sequence of the so-called strong homogeneous flows $\{\mathcal{C}_k^*\}$. We will show that it is asymptotically optimal for $\{\mathcal{P}_{S_k}\}$, yet it only yield the exact optimal solution if and only $k = 1, 2, 6, 10$. We note that $\{\mathcal{C}_k^*\}$ will also play important roles in terms of obtaining the exact optimal solutions; more specifically, as will be shown in Section 6, the optimal solution \mathcal{C}_k^{**} , $k = 3, 4, 5, 7, 8, 9$, are obtained using a perturbation from the corresponding \mathcal{C}_k^* .

4.1 Asymptotic Optimality

Definition 4.1. [Strong Homogeneous Flow] Let

$$\mathcal{C}_k^* := ((c_{i,j}^{*(1)}), (c_{i,j}^{*(2)}), \dots, (c_{i,j}^{*(k)})) \tag{11}$$

where

$$c_{i,j}^{*(\ell)} = \begin{cases} \frac{2}{k} - \frac{1}{k^2}, & \text{if } i = j = \ell, \\ \frac{1}{k} - \frac{1}{k^2}, & \text{if } i = \ell; j \neq \ell \text{ or } j = \ell; i \neq \ell, \\ -\frac{1}{k^2}, & \text{if } i \neq \ell \text{ and } j \neq \ell. \end{cases} \tag{12}$$

In the remainder of this paper, \mathcal{C}_k^* will be referred to as the *strong homogeneous flow*. Here we note that \mathcal{C}^* can be alternatively obtained by combining and concatenating elementary flows as in (IV.1) of [18].

Example 4.2. By definition, we have $\mathcal{C}_1^* = ((1))$ and

$$\mathcal{C}_2^* = \left(\left(\begin{pmatrix} \frac{3}{4} & \frac{1}{4} \\ \frac{1}{4} & \frac{1}{4} \end{pmatrix}, \begin{pmatrix} -\frac{1}{4} & \frac{1}{4} \\ \frac{1}{4} & \frac{3}{4} \end{pmatrix} \right) \right),$$

$$\mathcal{C}_3^* = \left(\left(\begin{pmatrix} \frac{5}{9} & \frac{2}{9} & \frac{2}{9} \\ \frac{2}{9} & \frac{-1}{9} & \frac{-1}{9} \\ \frac{2}{9} & \frac{-1}{9} & \frac{-1}{9} \end{pmatrix}, \begin{pmatrix} \frac{-1}{9} & \frac{2}{9} & \frac{-1}{9} \\ \frac{2}{9} & \frac{2}{9} & \frac{2}{9} \\ \frac{-1}{9} & \frac{2}{9} & \frac{-1}{9} \end{pmatrix}, \begin{pmatrix} \frac{-1}{9} & \frac{-1}{9} & \frac{2}{9} \\ \frac{-1}{9} & \frac{-1}{9} & \frac{-1}{9} \\ \frac{2}{9} & \frac{-1}{9} & \frac{2}{9} \end{pmatrix} \right).$$

Note that \mathcal{C}_2^* is the unique optimal point for $\mathcal{P}_{\mathcal{S}_2}$.

The following observation in [18] will serve as a key lemma in this paper.

Lemma 4.3 ([18]). *Let \mathcal{C}_k^* be the strong homogeneous multi-flow and \mathcal{S}_k be the strongly reachable sample set. Then, for all $s \in \mathcal{S}_k$,*

$$g_s(\mathcal{C}_k^*) = \frac{3k\alpha(s) - 2\beta(s)\alpha(s)}{k^2}. \quad (13)$$

We now define

$$\mathcal{S}_k(a, b) := \{s \in \mathcal{S}_k \mid \alpha(s) = a, \beta(s) = b\}.$$

The following two lemmas follow from Lemma 4.3 via straightforward computations.

Lemma 4.4. *For $\ell = 1, 2, \dots$, we have*

- 1) *If $k = 4\ell$, then $g_s(\mathcal{C}_k^*)$ reaches the maximum $\frac{9}{8}$ when $s \in \mathcal{S}_k(3\ell, 3\ell)$;*
- 2) *If $k = 4\ell + 1$, then $g_s(\mathcal{C}_k^*)$ reaches the maximum $\frac{18\ell^2 + 9\ell + 1}{16\ell^2 + 8\ell + 1}$ when $s \in \mathcal{S}_k(3\ell + 1, 3\ell + 1)$;*
- 3) *If $k = 4\ell + 2$, then $g_s(\mathcal{C}_k^*)$ reaches the maximum $\frac{9\ell^2 + 9\ell + 2}{8\ell^2 + 8\ell + 2}$ when $s \in \mathcal{S}_k(3\ell + 1, 3\ell + 1) \cup \mathcal{S}_k(3\ell + 2, 3\ell + 2)$;*
- 4) *If $k = 4\ell + 3$, then $g_s(\mathcal{C}_k^*)$ reaches the maximum $\frac{18\ell^2 + 27\ell + 10}{16\ell^2 + 24\ell + 9}$ when $s \in \mathcal{S}_k(3\ell + 2, 3\ell + 2)$.*

Lemma 4.5. *For $\ell = 1, 2, \dots$, we have*

- 1) *If $k = 4\ell$, then $g_s(\mathcal{C}_k^*)$ reaches the second largest value $\frac{9}{8} - \frac{2}{k^2}$ when $s \in \mathcal{S}_k(3\ell - 1, 3\ell - 1) \cup \mathcal{S}_k(3\ell + 1, 3\ell + 1)$;*
- 2) *If $k = 4\ell + 1$, then $g_s(\mathcal{C}_k^*)$ reaches the second largest value $\frac{18\ell^2 + 9\ell + 1}{16\ell^2 + 8\ell + 1} - \frac{1}{k^2}$ when $s \in \mathcal{S}_k(3\ell, 3\ell)$;*
- 3) *If $k = 4\ell + 2$, then $g_s(\mathcal{C}_k^*)$ reaches the second largest value $\frac{9\ell^2 + 9\ell + 2}{8\ell^2 + 8\ell + 2} - \frac{4}{k^2}$ when $s \in \mathcal{S}_k(3\ell, 3\ell) \cup \mathcal{S}_k(3\ell + 3, 3\ell + 3)$;*
- 4) *If $k = 4\ell + 3$, then $g_s(\mathcal{C}_k^*)$ reaches the second largest value $\frac{18\ell^2 + 27\ell + 10}{16\ell^2 + 24\ell + 9} - \frac{1}{k^2}$ when $s \in \mathcal{S}_k(3\ell + 3, 3\ell + 3)$.*

Definition 4.6. [**Asymptotically Optimal Solution**] A sequence $\{\mathcal{C}_k \mid \mathcal{C}_k \in \mathfrak{S}_k\}$ is said to be *asymptotically optimal* for $\{\mathcal{P}_{\mathcal{S}_k}\}$ if

$$\lim_{k \rightarrow \infty} g_{\mathcal{S}_k}(\mathcal{C}_k) = \lim_{k \rightarrow \infty} \mathcal{O}_{\mathcal{S}_k}.$$

The following theorem then immediately follows from Lemma 4.4:

Theorem 4.7. *$\{\mathcal{C}_k^*\}$ is asymptotically optimal for $\{\mathcal{P}_{\mathcal{S}_k}\}$.*

4.2 Optimality of \mathcal{C}_6^* and \mathcal{C}_{10}^*

In this section, we prove that \mathcal{C}_k^* is an optimal solution to \mathcal{P}_{S_k} if and only if $k = 1, 2, 6, 10$. We first state some needed notations and lemmas.

For any $\mathcal{C} \in \mathfrak{S}_k$, let $\mathcal{S}_k^\dagger(\mathcal{C})$ denote the set of all k -sample s such that

$$\begin{cases} g_s^{(\ell)}(\mathcal{C}) > 0, & \text{if } \ell \in \text{Ind}_s, \\ g_s^{(\ell)}(\mathcal{C}) < 0, & \text{if } \ell \notin \text{Ind}_s. \end{cases} \quad (14)$$

We then have the following lemma.

Lemma 4.8. *For any d , we have*

$$\mathcal{S}_k(d, d) \subset \mathcal{S}_k^\dagger(\mathcal{C}_k^*).$$

Proof. Notice that for each $s \in \mathcal{S}_k(d, d)$, $\alpha(s) = d < k$ and hence $g_s^{(\ell)}(\mathcal{C}_k^*)$ is the sum of at most $k - 1$ entries of $(c_{i,j}^{*(\ell)})$. By the definition of \mathcal{C}_k^* , we infer that if $\ell \in \text{Ind}_s$, then there exists at least one entry with value $(k - 1)/k^2$ or $(2k - 1)/k^2$ and the sum of the other entries are greater than or equal to $-(k - 2)/k^2$ and hence $g_s^{(\ell)}(\mathcal{C}_k^*) > 0$; and if $\ell \notin \text{Ind}_s$, then obviously $g_s^{(\ell)}(\mathcal{C}_k^*) = -d/k^2 < 0$. \square

An element in a k -sample s is said to be *diagonal* if its two coordinates are the same, otherwise *non-diagonal*. Let $\gamma(s)$ denote the number of diagonal elements in s . For example, let $s = \{(1, 1), (3, 3), (1, 2)(1, 4), (2, 5)\}$ be a 5-sample. Then, $(1, 1), (3, 3)$ are diagonal 5-samples, whereas $(1, 2), (1, 4), (2, 5)$ are non-diagonal 5-samples, and furthermore $\gamma(s) = 2$.

Lemma 4.9. *For any $s \in \mathcal{S}_k(d, d)$ with $d < k$, there exists a neighborhood $N(\mathcal{C}_k^*, \varepsilon) \subset \mathfrak{S}_k^{fix}$ of \mathcal{C}_k^* such that for all $\mathcal{C} \in N(\mathcal{C}_k^*, \varepsilon)$,*

$$g_s(\mathcal{C}) - g_s(\mathcal{C}_k^*) = \gamma(s)(\bar{x} + (2d - k - 1)\bar{y}) + (d - \gamma(s))((2\bar{a} + (2d - k - 2)\bar{b})), \quad (15)$$

where $\bar{a}, \bar{b}, \bar{x}, \bar{y}$ are defined by

$$\mathcal{C} - \mathcal{C}_k^* = \left(\left(\begin{array}{cccccc} \bar{x} & \bar{a} & \bar{a} & \dots & \bar{a} \\ \bar{a} & \bar{y} & \bar{b} & \dots & \bar{b} \\ \bar{a} & \bar{b} & \bar{y} & \dots & \bar{b} \\ \vdots & \vdots & & \ddots & \vdots \\ \bar{a} & \bar{b} & \bar{b} & \dots & \bar{y} \end{array} \right), \left(\begin{array}{cccccc} \bar{y} & \bar{a} & \bar{b} & \dots & \bar{b} \\ \bar{a} & \bar{x} & \bar{a} & \dots & \bar{a} \\ \bar{b} & \bar{a} & \bar{y} & \dots & \bar{b} \\ \vdots & \vdots & & \ddots & \vdots \\ \bar{b} & \bar{a} & \bar{b} & \dots & \bar{y} \end{array} \right), \dots, \left(\begin{array}{cccccc} \bar{y} & \bar{b} & \bar{b} & \dots & \bar{a} \\ \bar{b} & \bar{y} & \bar{b} & \dots & \bar{a} \\ \bar{b} & \bar{b} & \bar{y} & \dots & \bar{a} \\ \vdots & \vdots & & \ddots & \vdots \\ \bar{a} & \bar{a} & \bar{a} & \dots & \bar{x} \end{array} \right) \right).$$

Proof. Recall from Lemma 4.8 that $g_s^{(\ell)}(\mathcal{C}_k^*) > 0$ if $\ell \in \text{Ind}_s$ and $g_s^{(\ell)}(\mathcal{C}_k^*) < 0$ if $\ell \notin \text{Ind}_s$. Since each function $g_s^{(\ell)}(\cdot)$ is continuous, there exists a sufficiently small ε such that for all $\mathcal{C} \in N(\mathcal{C}_k^*, \varepsilon)$ and all $s \in \mathcal{S}_k(d, d)$, $g_s^{(\ell)}(\mathcal{C}) > 0$ if $\ell \in \text{Ind}_s$ and $g_s^{(\ell)}(\mathcal{C}) < 0$ if $\ell \notin \text{Ind}_s$. For any $\mathcal{C} \in N(\mathcal{C}_k^*, \varepsilon)$, we have

$$\begin{aligned} g_s(\mathcal{C}) - g_s(\mathcal{C}_k^*) &= \sum_{\ell=1}^k g_s^{(\ell)}(\mathcal{C}) - \sum_{\ell=1}^k g_s^{(\ell)}(\mathcal{C}_k^*) \\ &= \sum_{\ell \in \text{Ind}_s} (g_s^{(\ell)}(\mathcal{C}) - g_s^{(\ell)}(\mathcal{C}_k^*)) + \sum_{\ell \notin \text{Ind}_s} (g_s^{(\ell)}(\mathcal{C}_k^*) - g_s^{(\ell)}(\mathcal{C})). \end{aligned} \quad (16)$$

Noticing that $\alpha(s) = \beta(s) = d$, it is easy to check that

$$\sum_{\ell \in \text{Ind}_s} (g_s^{(\ell)}(\mathcal{C}) - g_s^{(\ell)}(\mathcal{C}_k^*)) = \gamma(s)\bar{x} + (d-1)\gamma(s)\bar{y} + 2(d-\gamma(s))\bar{a} + (d-2)(d-\gamma(s))\bar{b} \quad (17)$$

and

$$\sum_{\ell \notin \text{Ind}_s} (g_s^{(\ell)}(\mathcal{C}_k^*) - g_s^{(\ell)}(\mathcal{C})) = (d-k)(\gamma(s)\bar{y} + (d-\gamma(s))\bar{b}). \quad (18)$$

Combining Equations (17) and (18) and plugging the results into (16), we have

$$g_s(\mathcal{C}) - g_s(\mathcal{C}_k^*) = \gamma(s)(\bar{x} + (2d-k-1)\bar{y}) + (d-\gamma(s))(2\bar{a} + (2d-k-2)\bar{b}),$$

which completes the proof. \square

Lemma 4.10. *For any fixed $d < k$, there exists $\mathcal{C} \in \mathfrak{S}_k^{\text{fix}}$ such that $g_s(\mathcal{C}) - g_s(\mathcal{C}_k^*) < 0$ for all $s \in \mathcal{S}_k(d, d)$.*

Proof. Let $N(\mathcal{C}_k^*, \varepsilon)$ be the neighborhood of \mathcal{C}_k^* as in Lemma 4.9. We will prove that there exists $\mathcal{C} \in N(\mathcal{C}_k^*, \varepsilon)$ such that $g_s(\mathcal{C}) - g_s(\mathcal{C}_k^*) < 0$ for all $s \in \mathcal{S}_k(d, d)$. Note that, by Lemma 4.9, we only need to prove that there exist sufficiently small $\bar{a}, \bar{b}, \bar{x}, \bar{y}$ satisfying the following system:

$$\begin{cases} i(\bar{x} + (2d-k-1)\bar{y}) + (d-i)(2\bar{a} + (2d-k-2)\bar{b}), & i = 0, 1, \dots, d, \\ \bar{x} + (k-1)\bar{a} = 0, \\ \bar{y} + (k-2)\bar{b} + \bar{a} = 0. \end{cases}$$

Since the first and the $(d+1)$ -th inequalities imply the second to the d -th inequalities, we only need to prove there exist sufficiently small $\bar{a}, \bar{b}, \bar{x}, \bar{y}$ satisfying the following system:

$$\begin{cases} 2\bar{a} + (2d-k-2)\bar{b} < 0, \\ \bar{x} + (2d-k-1)\bar{y} < 0, \\ \bar{x} + (k-1)\bar{a} = 0, \\ \bar{y} + (k-2)\bar{b} + \bar{a} = 0, \end{cases} \quad (19)$$

or equivalently,

$$\begin{cases} 2\bar{a} + (2d-k-2)\bar{b} < 0, \\ 2\bar{a} + \frac{(2d-k-1)(k-2)}{d-1}\bar{b} > 0. \end{cases} \quad (20)$$

Since $d < k$, we have $\frac{2d-k-2}{2d-k-1} \neq \frac{k-2}{d-1}$ and hence $2d-k-2 \neq \frac{(2d-k-1)(k-2)}{d-1}$, which implies that there exist sufficiently small \bar{a} and \bar{b} such that (20) holds. By the last two equations of (19), \bar{x}, \bar{y} can also be chosen sufficiently small, which implies there exists $\mathcal{C} \in N(\mathcal{C}_k^*, \varepsilon)$ such that $g_s(\mathcal{C}) - g_s(\mathcal{C}_k^*) < 0$ for all $s \in \mathcal{S}_k(d, d)$, which completes the proof. \square

In what follows, a k -sample $s \in \mathcal{S}_k$ is said to be *maximizing* at \mathcal{C} if $g_s(\mathcal{C}) = g_{\mathcal{S}_k}(\mathcal{C})$, and we will use $\mathcal{S}_k^{\text{max}}(\mathcal{C})$ denote the set of all maximizing k -samples at \mathcal{C} . For a quick example, by 1) of Lemma 4.4, when $k = 4\ell$, any $s \in \mathcal{S}_k(3\ell, 3\ell)$ is a maximizing sample at \mathcal{C}_k^* ; and moreover, Lemma 4.4 implies that $\mathcal{S}_k(3\ell, 3\ell)$ is the set of all maximizing k -samples, i.e., $\mathcal{S}_k^{\text{max}}(\mathcal{C}_k^*) = \mathcal{S}_k(3\ell, 3\ell)$. Now, we are ready to give the main result of this section.

Theorem 4.11. \mathcal{C}_k^* is an optimal solution to $\mathcal{P}_{\mathcal{S}_k}$ if and only if $k = 1, 2, 6, 10$.

Proof. The case $k = 1$ is trivial and it is known [20] that \mathcal{C}_2^* is an optimal point for \mathcal{P}_2 . So we only need to prove \mathcal{C}_6^* and \mathcal{C}_{10}^* are respectively optimal points for \mathcal{P}_6 and \mathcal{P}_{10} , and \mathcal{C}_k^* is not an optimal point for $\mathcal{P}_{\mathcal{S}_k}$ when $k \neq 1, 2, 6, 10$.

In the remainder of the proof, we consider the following cases:

Case 1: $k > 1$ and $k \bmod 4 \neq 2$. In this case, by Lemma 4.4, there exists a d such that $\mathcal{S}_k^{max}(\mathcal{C}_k^*) = \mathcal{S}_k(d, d)$. Then, by Lemma 4.10, for some sufficiently small ε , we can choose $\mathcal{C} \in N(\mathcal{C}_k^*, \varepsilon)$ such that for any maximizing s at \mathcal{C}_k^* , the following two conditions hold: (1) $g_s(\mathcal{C}) < g_s(\mathcal{C}_k^*)$; (2) s is also a maximizing sample at \mathcal{C} (Here (2) is true because $|\mathcal{S}_k|$ is finite, and the function $g_s(\cdot)$ is continuous over \mathfrak{S}_k for each s). It then follows that $\max_{s \in \mathcal{S}_k} \{g_s(\mathcal{C})\} < \max_{s \in \mathcal{S}_k} \{g_s(\mathcal{C}_k^*)\}$, which means \mathcal{C}_k^* is not an optimal point for $\mathcal{P}_{\mathcal{S}_k}$.

Case 2: $k = 4\ell + 2$ for some integer $\ell \geq 3$. In this case, \mathcal{C}_k^* is not an optimal point for $\mathcal{P}_{\mathcal{S}_k}$. To prove this, we first show that for some sufficiently small ε , there exists $\mathcal{C} \in N(\mathcal{C}_k^*, \varepsilon)$ such that $g_s(\mathcal{C}) - g_s(\mathcal{C}_k^*) < 0$ for all $s \in \mathcal{S}_k(3\ell + 1, 3\ell + 1) \cup \mathcal{S}_k(3\ell + 2, 3\ell + 2)$ with $\ell \geq 3$. By Lemma 4.9, we only need to prove that there exist sufficiently small $\bar{a}, \bar{b}, \bar{x}, \bar{y}$ satisfying the following system:

$$\begin{cases} i(\bar{x} + (2(3\ell + 1) - k - 1)\bar{y} + (3\ell + 1 - i)(2\bar{a} + (2(3\ell + 1) - k - 2)\bar{b})) < 0, & i = 0, 1, \dots, 3\ell + 1, \\ i(\bar{x} + (2(3\ell + 2) - k - 1)\bar{y} + (3\ell + 2 - i)(2\bar{a} + (2(3\ell + 2) - k - 2)\bar{b})) < 0, & i = 0, 1, \dots, 3\ell + 2, \\ \bar{x} + (k - 1)\bar{a} = 0, \\ \bar{y} + (k - 2)\bar{b} + \bar{a} = 0. \end{cases} \quad (21)$$

Applying a similar argument as in the proof of Lemma 4.10 and noticing that $k = 4\ell + 2$, we simplify the above system to

$$\begin{cases} 2\bar{a} + 2(\ell - 1)\bar{b} < 0, \\ 2\bar{a} + 2\ell\bar{b} < 0, \\ 2\bar{a} + \frac{4(2\ell - 1)}{3}\bar{b} > 0, \\ 2\bar{a} + \frac{4\ell(2\ell + 1)}{3\ell + 1}\bar{b} > 0. \end{cases} \quad (22)$$

One then verifies that for all $\delta > 0$,

$$\begin{cases} \bar{a} = \frac{-11}{10}\ell\delta, \\ \bar{b} = \delta, \end{cases}$$

is a solution to (22). Choosing $\delta > 0$ small enough, we deduce that for some sufficiently small ε there exists $\mathcal{C} \in N(\mathcal{C}_k^*, \varepsilon)$ such that $g_s(\mathcal{C}) - g_s(\mathcal{C}_k^*) < 0$ for all $s \in \mathcal{S}_k(3\ell + 1, 3\ell + 1) \cup \mathcal{S}_k(3\ell + 2, 3\ell + 2)$ with $\ell \geq 3$. By the same reasoning as in Case 1, for some sufficiently small ε , we can choose $\mathcal{C} \in N(\mathcal{C}_k^*, \varepsilon)$ such that for any $s \in \mathcal{S}_k^{max}(\mathcal{C}_k^*)$, the following two hold: (1) $g_s(\mathcal{C}) < g_s(\mathcal{C}_k^*)$; (2) s is a maximizing sample of \mathcal{C} . Hence $\max_{s \in \mathcal{S}_k} \{g_s(\mathcal{C})\} < \max_{s \in \mathcal{S}_k} \{g_s(\mathcal{C}_k^*)\}$, which means \mathcal{C}_k^* is not an optimal point for $\mathcal{P}_{\mathcal{S}_k}$.

Case 3: $k = 4\ell + 2$ with $\ell = 1$, i.e., $k = 6$. In this case, consider Equation (22), which can be rewritten as

$$\begin{cases} \bar{a} < 0, \\ \bar{a} + \bar{b} < 0, \\ 3\bar{a} + 2\bar{b} > 0, \\ 2\bar{a} + 3\bar{b} > 0. \end{cases} \quad (23)$$

Note that this system has no solution because by the last two inequalities, we have $\bar{a} + \bar{b} > 0$, which contradicts the second inequality. Hence, within $N(\mathcal{C}_k^*, \varepsilon)$ (defined in Lemma 4.9), there is no point \mathcal{C} such that $g_s(\mathcal{C}) - g_s(\mathcal{C}_k^*) < 0$ for all maximizing s at \mathcal{C}_k^* (Note that by Lemma 4.4, the set of all such s is $\mathcal{S}_6^{max}(\mathcal{C}_6^*) = \mathcal{S}_6(4, 4) \cup \mathcal{S}_6(5, 5)$). Hence, \mathcal{C}_6^* is a local optimal point for $\mathcal{P}_{\mathcal{S}_6}$, and furthermore, by Lemma 2.2, \mathcal{C}_6^* is a global optimal point for $\mathcal{P}_{\mathcal{S}_6}$.

Case 4: $k = 4\ell + 2$ with $\ell = 2$, i.e., $k = 10$. In this case, consider Equation (22), which can be rewritten as

$$\begin{cases} \bar{a} + \bar{b} < 0, \\ \bar{a} + 2\bar{b} < 0, \\ \bar{a} + 2\bar{b} > 0, \\ 7\bar{a} + 20\bar{b} > 0. \end{cases} \quad (24)$$

Note that this system has no solution because the third inequality contradicts the second inequality. Hence, within $N(\mathcal{C}_k^*, \varepsilon)$ (defined in Lemma 4.9), there is no point \mathcal{C} such that $g_s(\mathcal{C}) - g_s(\mathcal{C}_k^*) < 0$ for all maximizing s at \mathcal{C}_k^* (Note that by Lemma 4.4, the set of all such s is $\mathcal{S}_{10}^{max}(\mathcal{C}_{10}^*) = \mathcal{S}_{10}(7, 7) \cup \mathcal{S}_{10}(8, 8)$). Hence, \mathcal{C}_{10}^* is a local optimal point for $\mathcal{P}_{\mathcal{S}_{10}}$, and again by Lemma 2.2, a global optimal point for $\mathcal{P}_{\mathcal{S}_{10}}$. \square

The following corollary says that the upper bound $\frac{9}{8}$ (derived in [20]) on $\mathcal{O}_{\mathcal{S}_k}$, $k \geq 3$, cannot be achieved.

Corollary 4.12. $\mathcal{O}_{\mathcal{S}_k} < \frac{9}{8}$ for all $k \geq 3$.

5 A Perturbation Framework

In this section, we propose a perturbation framework for the case $k \bmod 4 \neq 2$ that not only promises a better solution to $\mathcal{P}_{\mathcal{S}_k}$ than \mathcal{C}_k^* but also yields exact optimal solutions at least for some small k (see Section 6 for exact solutions for the cases $k = 3, 4, 5, 7, 8, 9$).

5.1 Valid Perturbation Direction

First of all, we define

$$L(\mathfrak{S}_k^{fix}) := \{\Delta = \mathcal{C} - \mathcal{C}' \mid \mathcal{C}, \mathcal{C}' \in \mathfrak{S}_k^{fix}\}.$$

Note that any $\Delta = (\Delta^{(1)}, \Delta^{(2)}, \dots, \Delta^{(k)}) \in L(\mathfrak{S}_k^{fix})$ can be written as

$$\Delta = \left(\left(\begin{array}{cccc} \bar{x} & \bar{a} & \bar{a} & \dots & \bar{a} \\ \bar{a} & \bar{y} & \bar{b} & \dots & \bar{b} \\ \bar{a} & \bar{b} & \bar{y} & \dots & \bar{b} \\ \vdots & \vdots & & \ddots & \vdots \\ \bar{a} & \bar{b} & \bar{b} & \dots & \bar{y} \end{array} \right), \left(\begin{array}{cccc} \bar{y} & \bar{a} & \bar{b} & \dots & \bar{b} \\ \bar{a} & \bar{x} & \bar{a} & \dots & \bar{a} \\ \bar{b} & \bar{a} & \bar{y} & \dots & \bar{b} \\ \vdots & \vdots & & \ddots & \vdots \\ \bar{b} & \bar{a} & \bar{b} & \dots & \bar{y} \end{array} \right), \dots, \left(\begin{array}{cccc} \bar{y} & \bar{b} & \bar{b} & \dots & \bar{a} \\ \bar{b} & \bar{y} & \bar{b} & \dots & \bar{a} \\ \bar{b} & \bar{b} & \bar{y} & \dots & \bar{a} \\ \vdots & \vdots & & \ddots & \vdots \\ \bar{a} & \bar{a} & \bar{a} & \dots & \bar{x} \end{array} \right) \right),$$

where $\bar{x}(\Delta) + (k-1)\bar{a}(\Delta) = 0$ and $\bar{y}(\Delta) + \bar{a}(\Delta) + (k-2)\bar{b}(\Delta) = 0$. Here, to emphasize the dependence, we have written $\bar{x}, \bar{y}, \bar{a}, \bar{b}$ as $\bar{x}(\Delta), \bar{y}(\Delta), \bar{a}(\Delta), \bar{b}(\Delta)$, respectively.

Lemma 5.1. For any fixed $d < k$, there exist $\Delta \in L(\mathfrak{S}_k^{fix})$ and $\varepsilon_0 > 0$ such that for all $s_1, s_2 \in \mathcal{S}_k(d, d)$ and all $0 < \varepsilon < \varepsilon_0$,

$$g_{s_1}(\mathcal{C}_k^* + \varepsilon\Delta) - g_{s_1}(\mathcal{C}_k^*) = g_{s_2}(\mathcal{C}_k^* + \varepsilon\Delta) - g_{s_2}(\mathcal{C}_k^*) < 0,$$

where $\varepsilon\Delta = (\varepsilon\Delta^{(1)}, \varepsilon\Delta^{(2)}, \dots, \varepsilon\Delta^{(k)})$.

Proof. To prove this lemma, we only need to slightly modify the proof of Lemma 4.10. More precisely, we assume $\bar{a} = (\frac{k^2}{2d} - k + 1)\bar{b}$, which implies $2\bar{a} + (2d - k - 2)\bar{b} = \bar{x} + (2d - k - 1)\bar{y}$, which is an extra constraint added to (19) ensuring (with the help of (15)) a uniform change from $g_s(\mathcal{C}_k^*)$ to $g_s(\mathcal{C}_k^* + \varepsilon\Delta)$ over all $s \in \mathcal{S}_k(d, d)$, i.e., for all $s_1, s_2 \in \mathcal{S}_k(d, d)$,

$$g_{s_1}(\mathcal{C}_k^* + \varepsilon\Delta) - g_{s_1}(\mathcal{C}_k^*) = g_{s_2}(\mathcal{C}_k^* + \varepsilon\Delta) - g_{s_2}(\mathcal{C}_k^*),$$

if ε is small enough. Moreover, it can be readily verified that the new system is still solvable with the extra constraint. Finally, choosing a solution and then \bar{a} (or equivalently, \bar{b}) properly similarly as in the proof of Lemma 4.10 yields the desired Δ . \square

Definition 5.2. [Valid Perturbation Direction] For any $k \bmod 4 \neq 2$, let d be such that the set of all maximizing k -samples at \mathcal{C}_k^* is $\mathcal{S}_k(d, d)$ (see Lemma 4.4). There is a unique *valid perturbation direction* $\Delta_k^* \in L(\mathfrak{S}_k^{fix})$ such that 1) $|\bar{a}(\Delta_k^*)| = 1$; 2) $\bar{a}(\Delta_k^*) = (\frac{k^2}{2d} - k + 1)\bar{b}(\Delta_k^*)$; 3) $2\bar{a}(\Delta_k^*) + (2d - k - 2)\bar{b}(\Delta_k^*) < 0$.

Remark 5.3. The ideas behind the above definition can be roughly explained as follows: As in the proof of Lemma 5.1, Conditions 2) and 3) will guarantee that the value of $g_s(\mathcal{C})$ uniformly decreases (over all $s \in \mathcal{S}_k(d, d)$) when perturbing \mathcal{C} from \mathcal{C}_k^* along the direction of Δ_k^* , and Condition 1) serves to “normalize” Δ_k^* to yield the uniqueness.

Remark 5.4. For $k \bmod 4 = 2$, there are two distinct d_1, d_2 such that the set of all maximizing k -samples is $\mathcal{S}_k(d_1, d_1) \cup \mathcal{S}_k(d_2, d_2)$ (see Lemma 4.4), and a perturbation direction that is valid with respect to $\mathcal{S}_k(d_1, d_1)$ may not be valid with respect to $\mathcal{S}_k(d_2, d_2)$. This is the key reason that our perturbation framework may not work for the case $k \bmod 4 = 2$, since it requires a uniform (over all $s \in \mathcal{S}_k^{max}(\mathcal{C}_k^*)$) decrease of the maximum in the course of perturbation.

Example 5.5. Let $k = 3$ and $d = 2$. It then follows from Condition (2) of Definition 5.2 that $\bar{b}(\Delta_3^*) = 4\bar{a}(\Delta_3^*)$. And by Condition (3), we infer that $\bar{a}(\Delta_3^*) > 0$. Moreover, by Condition (1), we have $\bar{a}(\Delta_3^*) = 1$, $\bar{b}(\Delta_3^*) = 4$, $\bar{x}(\Delta_3^*) = -2$ and $\bar{y}(\Delta_3^*) = -5$, or equivalently,

$$\Delta_3^* = \left(\left(\begin{pmatrix} -2 & 1 & 1 \\ 1 & -5 & 4 \\ 1 & 4 & -5 \end{pmatrix}, \begin{pmatrix} -5 & 1 & 4 \\ 1 & -2 & 1 \\ 4 & 1 & -5 \end{pmatrix}, \begin{pmatrix} -5 & 4 & 1 \\ 4 & -5 & 1 \\ 1 & 1 & -2 \end{pmatrix} \right) \right).$$

Similarly, set $k = 4$ and $d = 3$. Going through similar arguments as above, we have $\bar{a}(\Delta_4^*) = -1$, $\bar{b}(\Delta_4^*) = 3$, $\bar{x}(\Delta_4^*) = 3$ and $\bar{y}(\Delta_4^*) = -5$.

5.2 Valid Perturbation Size

In this section, assuming $k \bmod 4 \neq 2$, we discuss the valid perturbation size for Δ_k^* . For notational convenience, we will henceforth write

$$h_s^{(\ell)}(\varepsilon \Delta_k^*) := g_s^{(\ell)}(\mathcal{C}_k^* + \varepsilon \Delta_k^*) - g_s^{(\ell)}(\mathcal{C}_k^*), \quad h_s(\varepsilon \Delta_k^*) := g_s(\mathcal{C}_k^* + \varepsilon \Delta_k^*) - g_s(\mathcal{C}_k^*).$$

First of all, we need the following definition.

Definition 5.6. [Valid Perturbation Size] For a given k -sample s , $\varepsilon > 0$ is called $g_s^{(\ell)}$ -valid, $\ell = 1, 2, \dots, k$, if

$$g_s^{(\ell)}(\mathcal{C}_k^* + \varepsilon \Delta_k^*) \cdot g_s^{(\ell)}(\mathcal{C}_k^*) \geq 0;$$

and ε is called g_s -valid if for all $1 \leq \ell \leq k$, ε is $g_s^{(\ell)}$ -valid; ε is called $g_{\mathcal{S}_k}$ -valid if for all $s \in \mathcal{S}_k$, ε is g_s -valid.

Remark 5.7. Since the function $g_s^{(\ell)}$ is continuous, there always exists $\varepsilon > 0$ such that it is $g_s^{(\ell)}$ -valid, and furthermore, there always exists ε such that it is g_s -valid and $g_{\mathcal{S}_k}$ -valid.

We will also need the following two lemmas, whose proofs have been postponed to Appendices A and B, respectively.

Lemma 5.8. $\varepsilon > 0$ is $g_{\mathcal{S}_3}$ -valid if and only $\varepsilon \leq \frac{1}{36}$.

Lemma 5.9. $\varepsilon > 0$ is $g_{\mathcal{S}_4}$ -valid if and only $\varepsilon \leq \frac{1}{176}$.

5.3 Formula of $h_s(\varepsilon \Delta_k^*)$

In this section, assuming $k \bmod 4 \neq 2$, we will deduce a formula to compute $h_s(\varepsilon \Delta_k^*)$ for g_s -valid perturbations.

We start with the following definition.

Definition 5.10. [Type of a Sample] Let s_{diag} and s_{ndiag} denote the subsets of diagonal and non-diagonal elements of s , respectively. We say

$$\left\{ \begin{bmatrix} m_{Ind_{s_{diag}}}(1) \\ m_{Ind_{s_{ndiag}}}(1) \end{bmatrix}, \begin{bmatrix} m_{Ind_{s_{diag}}}(2) \\ m_{Ind_{s_{ndiag}}}(2) \end{bmatrix}, \dots, \begin{bmatrix} m_{Ind_{s_{diag}}}(k) \\ m_{Ind_{s_{ndiag}}}(k) \end{bmatrix} \right\} \quad (25)$$

is the *type* of s , which will be denoted by $T(s)$. And slightly abusing the notation, we may also use (25) to denote the set of all the samples of the type $T(s)$.

Example 5.11. For example, let $s = \{(1, 1), (3, 3), (1, 2)(1, 4), (2, 5)\}$ be a 5-sample. Then, $s_{diag} = \{(1, 1), (3, 3)\}$, $s_{ndiag} = \{(1, 2), (1, 4), (2, 5)\}$, and $T(s) = \left\{ \begin{bmatrix} 2 \\ 2 \end{bmatrix}, \begin{bmatrix} 0 \\ 2 \end{bmatrix}, \begin{bmatrix} 2 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right\}$.

Note that if two k -samples s_1, s_2 are in the same orbit of $Sym(k)$, namely, there exists $\sigma \in Sym(k)$ such that $s_1 = \sigma(s_2)$, then $T(s_1) = T(s_2)$, but the reverse direction does not hold in general. For example, one can check that $s_1 = \{(1, 1), (4, 3), (3, 4), (1, 5)\}$ and $s_2 = \{(1, 1), (1, 3), (3, 4), (4, 5)\}$ are not in the same orbit despite the fact they have the same type.

We have the following lemma, which says that for any $\mathcal{C} \in \mathfrak{S}_k^{fix}$, $g_s(\mathcal{C})$ is determined by $T(s)$. Note that the same statement may not hold true for $\mathcal{C} \notin \mathfrak{S}_k^{fix}$.

Lemma 5.12. Let $\mathcal{C} \in \mathfrak{S}_k^{fix}$. Then, for any k -samples s_1, s_2 with $T(s_1) = T(s_2)$, we have $g_{s_1}(\mathcal{C}) = g_{s_2}(\mathcal{C})$.

Proof. Since $T(s_1) = T(s_2)$, we can find a $\sigma \in \text{Sym}(k)$ such that for all $i = 1, 2, \dots, k$,

$$\begin{bmatrix} m_{\text{Ind}_{(s_1)_{diag}}}(i) \\ m_{\text{Ind}_{(s_1)_{ndiag}}}(i) \end{bmatrix} = \begin{bmatrix} m_{\text{Ind}_{(s_2)_{diag}}(\sigma(i))} \\ m_{\text{Ind}_{(s_2)_{ndiag}}(\sigma(i))} \end{bmatrix}.$$

Since $\mathcal{C} \in \mathfrak{S}_k^{fix}$, we have $g_{s_1}^{(i)}(\mathcal{C}) = g_{s_2}^{(\sigma(i))}(\mathcal{C})$. Hence,

$$g_{s_1}(\mathcal{C}) = \sum_{i=1}^k g_{s_1}^{(i)}(\mathcal{C}) = \sum_{i=1}^k g_{s_2}^{(\sigma(i))}(\mathcal{C}) = g_{s_2}(\mathcal{C}),$$

as desired. \square

We also need the following definition, which can be used to give an alternative classification of samples.

Definition 5.13. [**Discriminant**] For any k with $k \bmod 4 \neq 2$, the *discriminant* of a k -sample s is defined by

$$\mathcal{D}_k(s) := \bar{a}(\Delta_k^*) + \gamma(s)\bar{y}(\Delta_k^*) + (\alpha(s) - \gamma(s) - 1)\bar{b}(\Delta_k^*).$$

We next give an example for the above definition, for which we need to introduce more notation as follows: Let

$$\mathcal{S}_k(a, b, c) := \{s \in \mathcal{S}_k \mid \alpha(s) = a, \beta(s) = b, \gamma(s) = c\},$$

and

$$\mathcal{S}_k(a, b, c, d) := \{s \in \mathcal{S}_k \mid \alpha(s) = a, \beta(s) = b, \gamma(s) = c, \delta(s) = d\},$$

where

$$\delta(s) := |\{i \mid m_{\text{Ind}_s}(i) = 1\}|.$$

For example, one verifies that for

$$s = \{(1, 1), (1, 2), (2, 3), (3, 4), (6, 5)\},$$

we have $\text{Ind}_s = \{1, 1, 1, 2, 2, 3, 3, 4, 5, 6\}$, and moreover, $m_{\text{Ind}_s}(4) = m_{\text{Ind}_s}(5) = m_{\text{Ind}_s}(6) = 1$ and $\delta(s) = 3$, which imply that $s \in \mathcal{S}_6(5, 6, 1, 3)$.

Example 5.14. For the first case in Example 5.5, $\bar{x}(\Delta_3^*) = -2$, $\bar{y}(\Delta_3^*) = -5$, $\bar{a}(\Delta_3^*) = 1$ and $\bar{b}(\Delta_3^*) = 4$. Then, for any $s \in \mathcal{S}_3$,

$$\mathcal{D}_3(s) = 1 - 5\gamma(s) + 4(\alpha(s) - \gamma(s) - 1).$$

More specifically,

- if $s \in \mathcal{S}_3(3, 3, 0)$, then $\mathcal{D}_3(s) = 9$;

- if $s \in \mathcal{S}_3(3, 3, 1)$, then $\mathcal{D}_3(s) = 0$;
- if $s \in \mathcal{S}_3(3, 3, 2)$, then $\mathcal{D}_3(s) = -9$.

Similarly, for the second case in Example 5.5, $\bar{x}(\Delta_4^*) = 3$, $\bar{y}(\Delta_4^*) = -5$, $\bar{a}(\Delta_4^*) = -1$ and $\bar{b}(\Delta_4^*) = 3$. Hence, for any $s \in \mathcal{S}_4$,

$$\mathcal{D}_4(s) = -1 - 5\gamma(s) + 3(\alpha(s) - \gamma(s) - 1).$$

More specifically,

- if $s \in \mathcal{S}_4(4, 4, 0)$, then $\mathcal{D}_4(s) = 8$;
- if $s \in \mathcal{S}_4(4, 4, 1)$, then $\mathcal{D}_4(s) = 0$;
- if $s \in \mathcal{S}_4(4, 4, 2)$, then $\mathcal{D}_4(s) = -8$;
- if $s \in \mathcal{S}_4(4, 4, 3)$, then $\mathcal{D}_4(s) = -16$.

As will be shown below, the notion of discriminant can be used to give an alternative classification of samples.

Definition 5.15. [Class] A sample $s \in \mathcal{S}_k$ is said to be in class II if $s \in \mathcal{S}_k(k, k)$, $\delta(s) \neq 0$ and $\mathcal{D}_k(s) < 0$. Otherwise, it is said to be in class I.

Example 5.16. Using the fact $\mathcal{S}_3(3, 3, 2) = \mathcal{S}_3(3, 3, 2, 1)$ and recalling Example 5.14, we have that $s \in \mathcal{S}_3$ is in class II if and only if $s \in \mathcal{S}_3(3, 3, 2)$. Similarly, we have that $s \in \mathcal{S}_4$ is in class II if and only if $s \in \mathcal{S}_4(4, 4, 2) \cup \mathcal{S}_4(4, 4, 3)$ and $\delta(s) \neq 0$. It is easy to verify that

$$\begin{aligned} \mathcal{S}_4(4, 4, 2) = & \left\{ \begin{bmatrix} 2 \\ 2 \end{bmatrix}, \begin{bmatrix} 2 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right\} \cup \left\{ \begin{bmatrix} 2 \\ 1 \end{bmatrix}, \begin{bmatrix} 2 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right\} \\ & \cup \left\{ \begin{bmatrix} 2 \\ 1 \end{bmatrix}, \begin{bmatrix} 2 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 2 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right\} \cup \left\{ \begin{bmatrix} 2 \\ 0 \end{bmatrix}, \begin{bmatrix} 2 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 2 \end{bmatrix}, \begin{bmatrix} 0 \\ 2 \end{bmatrix} \right\}, \end{aligned}$$

where $\delta(s) = 0$ if and only if $s \in \left\{ \begin{bmatrix} 2 \\ 0 \end{bmatrix}, \begin{bmatrix} 2 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 2 \end{bmatrix}, \begin{bmatrix} 0 \\ 2 \end{bmatrix} \right\}$. This, together with the fact that $\mathcal{S}_4(4, 4, 3) = \mathcal{S}_4(4, 4, 3, 1)$, implies that all the class II samples of \mathcal{S}_4 are

$$\left\{ \begin{bmatrix} 2 \\ 2 \end{bmatrix}, \begin{bmatrix} 2 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right\} \cup \left\{ \begin{bmatrix} 2 \\ 1 \end{bmatrix}, \begin{bmatrix} 2 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right\} \cup \left\{ \begin{bmatrix} 2 \\ 1 \end{bmatrix}, \begin{bmatrix} 2 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 2 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right\} \cup \mathcal{S}_4(4, 4, 3).$$

The following lemma, which is the main result of this section, measures how much $g_s(\cdot)$ changes from \mathcal{C}_k^* under a valid perturbation along the direction of Δ_k^* .

Lemma 5.17. *Let $s \in \mathcal{S}_k$ and $\varepsilon > 0$ be g_s -valid. Then,*

$$h_s(\varepsilon \Delta_k^*) = A_s \cdot \bar{x}(\varepsilon \Delta_k^*) + B_s \cdot \bar{y}(\varepsilon \Delta_k^*) + C_s \cdot \bar{a}(\varepsilon \Delta_k^*) + D_s \cdot \bar{b}(\varepsilon \Delta_k^*), \quad (26)$$

where, if s is in class I, then

$$\begin{aligned} A_s &= \gamma(s), \\ B_s &= \gamma(s)(2\beta(s) - k - 1), \\ C_s &= 2(\alpha(s) - \gamma(s)), \\ D_s &= (\alpha(s) - \gamma(s))(2\beta(s) - k - 2); \end{aligned} \quad (27)$$

and if s is in class II, then

$$\begin{aligned}
A_s &= \gamma(s), \\
B_s &= \gamma(s)(k - 2\delta(s) - 1), \\
C_s &= 2(k - \gamma(s) - \delta(s)), \\
D_s &= k^2 - (2\delta(s) + \gamma(s) + 2)k + 2\gamma(s)\delta(s) + 2\gamma(s) + 2\delta(s).
\end{aligned} \tag{28}$$

Proof. For any $s \in \mathcal{S}_k$ and any $1 \leq \ell \leq k$, by the definition of \mathcal{C}_k^* , it is easy to see that

$$g_s^{(\ell)}(\mathcal{C}_k^*) = \frac{m_{Ind_s}(\ell)}{k} - \frac{\alpha(s)}{k^2}, \tag{29}$$

which immediately implies that

- $g_s^{(\ell)}(\mathcal{C}_k^*) = 0$ if $\alpha(s) = k$ and $m_{Ind_s}(\ell) = 1$;
- $g_s^{(\ell)}(\mathcal{C}_k^*) < 0$ if $m_{Ind_s}(\ell) = 0$, i.e., $\ell \notin Ind_s$;
- $g_s^{(\ell)}(\mathcal{C}_k^*) > 0$ otherwise.

We first consider the samples in class I. By definition, there are the following three cases: (1) $s \notin \mathcal{S}_k(k, k)$; (2) $s \in \mathcal{S}_k(k, k)$ and $\delta(s) = 0$; (3) $s \in \mathcal{S}_k(k, k)$, $\delta(s) \neq 0$ and $\mathcal{D}_k(s) \geq 0$.

By the above discussions, for Cases (1) and (2), we have $g_s^{(\ell)}(\mathcal{C}_k^*) \neq 0$ for all $1 \leq \ell \leq k$. Then, by the definition of a valid perturbation, the following hold for the g_s -valid ε :

$$\begin{aligned}
&\text{if } g_s^{(\ell)}(\mathcal{C}_k^*) > 0, \text{ then } g_s^{(\ell)}(\mathcal{C}_k^* + \varepsilon\Delta_k^*) \geq 0; \\
&\text{if } g_s^{(\ell)}(\mathcal{C}_k^*) < 0, \text{ then } g_s^{(\ell)}(\mathcal{C}_k^* + \varepsilon\Delta_k^*) \leq 0.
\end{aligned} \tag{30}$$

For Case (3), since ε is g_s -valid, (30) still holds. In this case, since $\delta(s) \neq 0$, there exists some ℓ such that $g_s^{(\ell)}(\mathcal{C}_k^*) = 0$; and for such an ℓ , it can be verified that

$$g_s^{(\ell)}(\mathcal{C}_k^* + \varepsilon\Delta_k^*) = \bar{a}(\varepsilon\Delta_k^*) + \gamma(s)\bar{y}(\varepsilon\Delta_k^*) + (\alpha(s) - \gamma(s) - 1)\bar{b}(\varepsilon\Delta_k^*) = \mathcal{D}_k(s) \geq 0. \tag{31}$$

Now, combining (30) and (31), we deduce that

$$\begin{aligned}
&\text{if } \ell \in Ind_s, \text{ then } g_s^{(\ell)}(\mathcal{C}_k^*) \geq 0 \text{ and } g_s^{(\ell)}(\mathcal{C}_k^* + \varepsilon\Delta_k^*) \geq 0; \\
&\text{if } \ell \notin Ind_s, \text{ then } g_s^{(\ell)}(\mathcal{C}_k^*) < 0 \text{ and } g_s^{(\ell)}(\mathcal{C}_k^* + \varepsilon\Delta_k^*) \leq 0.
\end{aligned} \tag{32}$$

Hence, for any sample s in class I, we have

$$\begin{aligned}
h_s(\varepsilon\Delta_k^*) &= \sum_{\ell=1}^k |g_s^{(\ell)}(\mathcal{C}_k^* + \varepsilon\Delta_k^*)| - \sum_{\ell=1}^k |g_s^{(\ell)}(\mathcal{C}_k^*)| \\
&= \sum_{\ell \in Ind_s} (g_s^{(\ell)}(\mathcal{C}_k^* + \varepsilon\Delta_k^*) - g_s^{(\ell)}(\mathcal{C}_k^*)) - \sum_{\ell \notin Ind_s} (g_s^{(\ell)}(\mathcal{C}_k^* + \varepsilon\Delta_k^*) - g_s^{(\ell)}(\mathcal{C}_k^*)) \\
&= \sum_{\ell \in Ind_s} h_s^{(\ell)}(\varepsilon\Delta_k^*) - \sum_{\ell \notin Ind_s} h_s^{(\ell)}(\varepsilon\Delta_k^*).
\end{aligned} \tag{33}$$

Note that

$$\begin{aligned} \sum_{\ell \in \text{Ind}_s} h_s^{(\ell)}(\varepsilon \Delta_k^*) &= \gamma(s) \bar{x}(\varepsilon \Delta_k^*) + (\beta(s) - 1) \gamma(s) \bar{y}(\varepsilon \Delta_k^*) \\ &\quad + 2(\alpha(s) - \gamma(s)) \bar{a}(\varepsilon \Delta_k^*) + (\alpha(s) - \gamma(s)) (\beta(s) - 2) \gamma(s) \bar{b}(\varepsilon \Delta_k^*) \end{aligned}$$

and

$$\sum_{\ell \notin \text{Ind}_s} h_s^{(\ell)}(\varepsilon \Delta_k^*) = (k - \beta(s)) (\gamma(s) \bar{y}(\varepsilon \Delta_k^*) + (\alpha(s) - \gamma(s)) \bar{b}(\varepsilon \Delta_k^*)).$$

Substituting the above equalities into (33) then yields the result for class I.

Now, we consider the samples in class II. By definition, there exists some ℓ such that $g_s^{(\ell)}(\mathcal{C}_k^*) = 0$; and for such an ℓ ,

$$g_s^{(\ell)}(\mathcal{C}_k^* + \varepsilon \Delta_k^*) = \bar{a}(\Delta_k^*) + \gamma(s) \bar{y}(\Delta_k^*) + (\alpha(s) - \gamma(s) - 1) \bar{b}(\Delta_k^*) = \mathcal{D}_k(s) < 0.$$

Hence, similarly as above, we have

$$\begin{aligned} h_s(\varepsilon \Delta_k^*) &= \sum_{\ell=1}^k |g_s^{(\ell)}(\mathcal{C}_k^* + \varepsilon \Delta_k^*)| - \sum_{\ell=1}^k |h_s^{(\ell)}(\mathcal{C}_k^*)| \\ &= \sum_{\ell \in \text{Ind}_s} h_s^{(\ell)}(\varepsilon \Delta_k^*) - \sum_{\ell \notin \text{Ind}_s} h_s^{(\ell)}(\varepsilon \Delta_k^*) - 2\delta(s) \mathcal{D}_k(s). \end{aligned} \tag{34}$$

Noting that $\alpha(s) = \beta(s) = k$ for any class II sample s and substituting for the values of $\sum_{\ell \in \text{Ind}_s} h_s^{(\ell)}(\varepsilon \Delta_k^*)$, $\sum_{\ell \notin \text{Ind}_s} h_s^{(\ell)}(\varepsilon \Delta_k^*)$ as in the proof for class I, the result for class II then follows, which completes the proof. \square

5.4 Perturbation framework

Note that by Theorem 4.11, for any $k \neq 1, 2, 6, 10$, one can perturb \mathcal{C}_k^* to obtain a better solution to $\mathcal{P}_{\mathcal{S}_k}$, which however may not be optimal. In the following, we propose a framework of perturbing \mathcal{C}_k^* to obtain \mathcal{C}_k^{**} for $k \bmod \neq 2$, which are optimal at least for the cases $k = 3, 4, 5, 7, 9$ (see Section 6).

Step 1: Compute Δ_k^* . This step can be done by solving 1), 2) and 3) in Definition 5.2.

Step 2: Compute \mathcal{C}_k^{} .** For this step, we first use Lemmas 4.4 and 4.5 to obtain the subsets of samples which achieves the maximum and the second largest values of $\{g_s(\mathcal{C}_k^*) \mid s \in \mathcal{S}_k\}$. And we then use Lemma 5.17 to compute $h_s(\varepsilon \Delta_k^*)$ for all $s \in \mathcal{S}_k$. In the end, we increase the value of ε from 0 so that the maximum will decrease (uniformly over all $s \in \mathcal{S}_k^{max}(\mathcal{C}_k^*)$) until it meets the increasing second largest value at $\varepsilon = \varepsilon^*$, and then set $\mathcal{C}_k^{**} = \mathcal{C}_k^* + \varepsilon^* \Delta_k^*$.

Step 3: Compute $\mathcal{S}_k^{max}(\mathcal{C}_k^{})$.** We first check by Definition 5.6 the validity of ε obtained in **Step 2**. It turns out that for each k , there might exist a small number of samples s for which ε is not g_s -valid. For such s , we can simply compute the value of $g_s(\mathcal{C}_k^{**})$ using the definition of g_s , and then we compute, by using Lemma 4.3 and Lemma 5.17, the value of $g_s(\mathcal{C}_k^{**}) = g_s(\mathcal{C}_k^*) + h_s(\varepsilon \Delta_k^*)$ for all s where ε is g_s -valid. Finally, with the values of all $g_s(\mathcal{C}_k^{**})$, we derive $\mathcal{S}_k^{max}(\mathcal{C}_k^{**})$.

6 Optimal Solutions for $k = 3, 4, 5, 7, 8, 9$

In this section, through perturbing the corresponding \mathcal{C}_k^* , we obtain the optimal solutions \mathcal{C}_k^{**} to $\mathcal{P}_{\mathcal{S}_k}$ for $k = 3, 4, 5, 7, 8, 9$, and we further establish the uniqueness of these optimal solutions.

6.1 From \mathcal{C}_k^* to \mathcal{C}_k^{**} for $k = 3, 4, 5, 7, 9$

The perturbation from \mathcal{C}_k^* to \mathcal{C}_k^{**} follows from the framework in Section 5.4 with however some possible simplifications and adaptations to varying degrees for different k .

■ We first deal with the case $k = 3$ through the following steps.

Step 1: Compute Δ_3^* . This has already been done in Example 5.5.

Step 2: Compute \mathcal{C}_3^{} .** For this step, we need to compute $h_s(\varepsilon\Delta_3^*)$ for all $s \in \mathcal{S}_3$. To this end, we compute using Lemma 5.17,

$$h_s(\varepsilon\Delta_3^*) = (-2A_s - 5B_s + C_s + 4D_s)\varepsilon.$$

By Example 5.16, s is in class II if and only if $s \in \mathcal{S}_3(3, 3, 2, 1) = \mathcal{S}_3(3, 3, 2)$. Then, by (28), we have $A_s = 2$, $B_s = C_s = 0$, $D_s = 1$ and hence $h_s(\varepsilon\Delta_3^*) = 0$ for $s \in \mathcal{S}_3(3, 3, 2)$. For any sample s in class I, we use (27) to compute the coefficients and then compute $h_s(\varepsilon\Delta_3^*)$. The computations as above yield Table 1, where the values of all $g_s(\mathcal{C}_3^*)$ and $h_s(\varepsilon\Delta_3^*)$ are listed.

Table 1: The values of $g_s(\mathcal{C}_3^*)$ and $h_s(\varepsilon\Delta_3^*)$

Class of s	class I			
Subclass of s	$\mathcal{S}_3(1, 1, 1)$	$\mathcal{S}_3(1, 2, 0)$	$\mathcal{S}_3(2, 2)$	$\mathcal{S}_3(2, 3, 0)$
$g_s(\mathcal{C}_3^*)$	$\frac{7}{9}$	$\frac{5}{9}$	$\frac{10}{9}$	$\frac{6}{9}$
$h_s(\varepsilon\Delta_3^*)$	8ε	-2ε	-4ε	12ε

Class of s	class I				class II
Subclass of s	$\mathcal{S}_3(2, 3, 1)$	$\mathcal{S}_3(3, 3, 0)$	$\mathcal{S}_3(3, 3, 1)$	$\mathcal{S}_3(3, 3, 3)$	$\mathcal{S}_3(3, 3, 2)$
$g_s(\mathcal{C}_3^*)$	$\frac{6}{9}$	1			
$h_s(\varepsilon\Delta_3^*)$	-6ε	18ε	0	-36ε	0

By Table 1, $g_s(\mathcal{C}_3^*)$ achieves the maximum $\frac{10}{9}$ at $\mathcal{S}_3(2, 2)$ (or, more precisely, at any sample from $\mathcal{S}_3(2, 2)$) and the second largest value 1 at $\mathcal{S}_3(3, 3)$. Now, we will perturb \mathcal{C}_k^* along the direction of Δ_k^* to obtain \mathcal{C}_k^{**} so that, roughly speaking, the maximum will decrease until it meets the increasing second largest value. To this end, we note that in the course of perturbation, the second largest value is always achieved at $\mathcal{S}_3(3, 3)$, and we thereby solve $1 + 18\varepsilon = \frac{10}{9} - 4\varepsilon$, which yields $\varepsilon^* = \frac{1}{22 \times 9} = \frac{1}{198}$ and furthermore,

$$\mathcal{C}_3^{**} := \Delta_3^* \times \frac{1}{198} + \mathcal{C}_3^* = \left(\left(\begin{pmatrix} \frac{12}{22} & \frac{5}{22} & \frac{5}{22} \\ \frac{5}{22} & \frac{22}{22} & \frac{22}{22} \\ \frac{5}{22} & \frac{22}{22} & \frac{22}{22} \end{pmatrix}, \begin{pmatrix} \frac{-3}{22} & \frac{5}{22} & \frac{-2}{22} \\ \frac{5}{22} & \frac{22}{22} & \frac{22}{22} \\ \frac{-2}{22} & \frac{5}{22} & \frac{-3}{22} \end{pmatrix}, \begin{pmatrix} \frac{-3}{22} & \frac{-2}{22} & \frac{5}{22} \\ \frac{22}{22} & \frac{22}{22} & \frac{22}{22} \\ \frac{5}{22} & \frac{5}{22} & \frac{12}{22} \end{pmatrix} \right).$$

By Lemma 5.8, $\Delta_3^* \times \frac{1}{198}$ is a valid perturbation.

Step 3: Compute $\mathcal{S}_3^{max}(\mathcal{C}_3^{})$.** From Table 1, it is easy to verify that $\{g_s(\mathcal{C}_3^{**}) \mid s \in \mathcal{S}_3\}$ achieves the maximum $\frac{12}{11}$ at $\mathcal{S}_3(2, 2) \cup \mathcal{S}_3(3, 3, 0)$. In other words, $\mathcal{S}_k^{max}(\mathcal{C}_3^{**}) = \mathcal{S}_3(2, 2) \cup \mathcal{S}_3(3, 3, 0)$.

■ Now, we focus on the case $k = 4$ through the following steps.

Step 1: Compute Δ_4^* . This has already been done in Example 5.5.

Step 2: Compute \mathcal{C}_4^{} .** For this step, we need to compute $h_s(\varepsilon\Delta_4^*)$ for all $s \in \mathcal{S}_4$. To this end, we compute using Lemma 5.17,

$$h_s(\varepsilon\Delta_4^*) = (3A_s - 5B_s - C_s + 3D_s)\varepsilon.$$

By Example 5.16, $s \in \mathcal{S}_4$ is in class II if and only if

$$s \in \left\{ \begin{bmatrix} 2 \\ 2 \end{bmatrix}, \begin{bmatrix} 2 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right\} \cup \left\{ \begin{bmatrix} 2 \\ 1 \end{bmatrix}, \begin{bmatrix} 2 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right\} \cup \left\{ \begin{bmatrix} 2 \\ 1 \end{bmatrix}, \begin{bmatrix} 2 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 2 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right\} \cup \mathcal{S}_4(4, 4, 3).$$

For the class II samples, if $s \in \left\{ \begin{bmatrix} 2 \\ 2 \end{bmatrix}, \begin{bmatrix} 2 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right\} \cup \left\{ \begin{bmatrix} 2 \\ 1 \end{bmatrix}, \begin{bmatrix} 2 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right\}$, we have $\gamma(s) = \delta(s) = 2$ and by (28) $A_s = 2, B_s = -2, C_s = D_s = 0$ and hence $h_s(\varepsilon\Delta_4^*) = 16\varepsilon$; if $s \in \left\{ \begin{bmatrix} 2 \\ 1 \end{bmatrix}, \begin{bmatrix} 2 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 2 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right\}$, we have $\gamma(s) = 2, \delta(s) = 1$ and by (28) $A_s = B_s = C_s = D_s = 2$ and hence $h_s(\varepsilon\Delta_4^*) = 0$; if $s \in \mathcal{S}_4(4, 4, 3)$, we have $\gamma(s) = 3, \delta(s) = 1$ and by (28) $A_s = B_s = 3, C_s = 0$ and $D_s = 2$ and hence $h_s(\varepsilon\Delta_4^*) = 0$. For the samples in class I, we use (27) to compute the coefficients and then obtain $h_s(\varepsilon\Delta_4^*)$. The computations as above yield Table 2, where the values of all $g_s(\mathcal{C}_4^*)$ and $h_s(\varepsilon\Delta_4^*)$ are listed.

Table 2: The values of $g_s(\mathcal{C}_4^*)$ and $h_s(\varepsilon\Delta_4^*)$. Note that $\mathcal{S}_4(2, 3) = \mathcal{S}_4(2, 3, 0) \cup \mathcal{S}_4(2, 3, 1)$.

Class of s	class I				
Subclass of s	$\mathcal{S}_4(1, 1, 1)$	$\mathcal{S}_4(1, 2, 0)$	$\mathcal{S}_4(2, 2, 0)$	$\mathcal{S}_4(2, 2, 1)$	$\mathcal{S}_4(2, 2, 2)$
$g_s(\mathcal{C}_4^*)$	$\frac{5}{8}$	$\frac{1}{2}$	1		
$h_s(\varepsilon\Delta_4^*)$	18ε	-8ε	-16ε	0	16ε

Class of s	class I					
Subclass of s	$\mathcal{S}_4(2, 3)$	$\mathcal{S}_4(2, 4, 0)$	$\mathcal{S}_4(3, 3)$	$\mathcal{S}_4(3, 4, 0)$	$\mathcal{S}_4(3, 4, 1)$	$\mathcal{S}_4(3, 4, 2)$
$g_s(\mathcal{C}_4^*)$	$\frac{3}{4}$	$\frac{1}{2}$	$\frac{9}{8}$	$\frac{3}{4}$		
$h_s(\varepsilon\Delta_4^*)$	-4ε	8ε	-6ε	12ε	-4ε	-20ε

Class of s	class I			class II
Subclass of s	$\mathcal{S}_4(4, 4, 0)$	$\mathcal{S}_4(4, 4, 1)$	$\left\{ \begin{bmatrix} 2 \\ 0 \end{bmatrix}, \begin{bmatrix} 2 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 2 \end{bmatrix}, \begin{bmatrix} 0 \\ 2 \end{bmatrix} \right\}$	$\left\{ \begin{bmatrix} 2 \\ 1 \end{bmatrix}, \begin{bmatrix} 2 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 2 \end{bmatrix} \right\}$
$g_s(\mathcal{C}_4^*)$	1			
$h_s(\varepsilon\Delta_4^*)$	16ε	0	-16ε	0

Class of s	class I	class II	
Subclass of s	$\mathcal{S}_4(4, 4, 4)$	$\left\{ \begin{bmatrix} 2 \\ 1 \end{bmatrix}, \begin{bmatrix} 2 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right\}$	$\left\{ \begin{bmatrix} 2 \\ 2 \end{bmatrix}, \begin{bmatrix} 2 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right\} \cup \mathcal{S}_4(4, 4, 3)$
$g_s(\mathcal{C}_4^*)$	1		
$h_s(\varepsilon\Delta_4^*)$	-48ε	16ε	0

Note that by Table 2, $\{g_s(\mathcal{C}_4^*) \mid s \in \mathcal{S}_k\}$ achieves the maximum $\frac{9}{8}$ at $\mathcal{S}_4(3, 3)$ and the second largest value 1 at $\mathcal{S}_4(2, 2) \cup \mathcal{S}_4(4, 4)$. Now, similarly as in the case $k = 3$, we will perturb

\mathcal{C}_4^* along the direction of Δ_4^* to obtain \mathcal{C}_4^{**} . To this end, we again note that in the course of perturbation, the second largest value is always achieved at $\mathcal{S}_4(4, 4)$, and we thereby solve $1 + 16\varepsilon = \frac{9}{8} - 6\varepsilon$, which yields $\varepsilon^* = \frac{1}{22 \times 8} = \frac{1}{176}$, and furthermore,

$$\begin{aligned} \mathcal{C}_4^{**} &= \Delta_4^* \times \frac{1}{176} + \mathcal{C}_4^* \\ &= \left(\left(\begin{pmatrix} \frac{5}{11} & \frac{2}{11} & \frac{2}{11} & \frac{2}{11} \\ \frac{2}{11} & \frac{2}{11} & \frac{2}{11} & \frac{2}{11} \\ \frac{2}{11} & \frac{2}{11} & \frac{2}{11} & \frac{2}{11} \\ \frac{2}{11} & \frac{2}{11} & \frac{2}{11} & \frac{2}{11} \end{pmatrix}, \begin{pmatrix} \frac{-1}{11} & \frac{2}{11} & \frac{-1}{11} & \frac{-1}{11} \\ \frac{2}{11} & \frac{2}{11} & \frac{2}{11} & \frac{2}{11} \\ \frac{-1}{11} & \frac{2}{11} & \frac{-1}{11} & \frac{-1}{11} \\ \frac{-1}{11} & \frac{2}{11} & \frac{-1}{11} & \frac{-1}{11} \end{pmatrix}, \begin{pmatrix} \frac{-1}{11} & \frac{-1}{11} & \frac{2}{11} & \frac{-1}{11} \\ \frac{-1}{11} & \frac{-1}{11} & \frac{2}{11} & \frac{-1}{11} \\ \frac{2}{11} & \frac{2}{11} & \frac{2}{11} & \frac{2}{11} \\ \frac{-1}{11} & \frac{-1}{11} & \frac{2}{11} & \frac{-1}{11} \end{pmatrix}, \begin{pmatrix} \frac{-1}{11} & \frac{-1}{11} & \frac{-1}{11} & \frac{2}{11} \\ \frac{-1}{11} & \frac{-1}{11} & \frac{-1}{11} & \frac{2}{11} \\ \frac{-1}{11} & \frac{-1}{11} & \frac{-1}{11} & \frac{2}{11} \\ \frac{-1}{11} & \frac{-1}{11} & \frac{-1}{11} & \frac{2}{11} \end{pmatrix} \right). \end{aligned}$$

By Lemma 5.9, $\Delta_4^* \times \frac{1}{176}$ is a valid perturbation.

Step 3: Compute $\mathcal{S}_4^{max}(\mathcal{C}_4^{})$.** From Table 2, it is easy to verify that $\{g_s(\mathcal{C}_4^{**}) \mid s \in \mathcal{S}_4\}$ achieves the maximum $\frac{12}{11}$ at

$$\mathcal{S}_4^{max}(\mathcal{C}_4^{**}) = \mathcal{S}_4(2, 2, 2) \cup \mathcal{S}_4(3, 3) \cup \mathcal{S}_4(4, 4, 0) \cup \left\{ \begin{bmatrix} 2 \\ 1 \end{bmatrix}, \begin{bmatrix} 2 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right\} \cup \left\{ \begin{bmatrix} 2 \\ 2 \end{bmatrix}, \begin{bmatrix} 2 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right\}.$$

■ For the cases $k = 5, 7, 8, 9$, we only outline the major steps to derive \mathcal{C}_k^{**} without giving all the computation details.

Step 1: Compute Δ_k^* .

- For $k = 5$, $\bar{x}(\Delta_5^*) = 4$, $\bar{a}(\Delta_5^*) = -1$, $\bar{b}(\Delta_5^*) = \frac{8}{7}$, $\bar{y}(\Delta_5^*) = -\frac{17}{7}$.
- For $k = 7$, $\bar{x}(\Delta_7^*) = 6$, $\bar{a}(\Delta_7^*) = -1$, $\bar{b}(\Delta_7^*) = \frac{10}{11}$, $\bar{y}(\Delta_7^*) = -\frac{39}{11}$.
- For $k = 8$, $\bar{x}(\Delta_8^*) = 7$, $\bar{a}(\Delta_8^*) = -1$, $\bar{b}(\Delta_8^*) = \frac{3}{5}$, $\bar{y}(\Delta_8^*) = -\frac{13}{5}$.
- For $k = 9$, $\bar{x}(\Delta_9^*) = 8$, $\bar{a}(\Delta_9^*) = -1$, $\bar{b}(\Delta_9^*) = \frac{14}{31}$, $\bar{y}(\Delta_9^*) = -\frac{67}{31}$.

Step 2: Compute \mathcal{C}_k^{} .**

- For $k = 5$, $g_s(\mathcal{C}_k^*)$ achieves the maximum $\frac{28}{25}$ at $\mathcal{S}_5(4, 4)$ and the second largest value $\frac{27}{25}$ at $\mathcal{S}_5(3, 3)$. By Definition 5.15, all these samples are of class I. Then, an application of Lemma 5.17 yields that

$$h_s(\varepsilon \Delta_5^*) = \begin{cases} -\frac{24}{7}\varepsilon, & s \in \mathcal{S}_5(4, 4), \\ \frac{50\gamma(s)-66}{7}\varepsilon, & s \in \mathcal{S}_5(3, 3), \end{cases}$$

based on which, we infer that the second largest value increases the fastest (with speed $h_s(\varepsilon \Delta_5^*) = \frac{84\varepsilon}{7}$) when $\gamma(s) = 3$. Solving the equation $\frac{28}{25} - \frac{24}{7}\varepsilon = \frac{27}{25} + \frac{84}{7}\varepsilon$, we have $\varepsilon^* = \frac{7}{108 \times 25}$ and obtain $\mathcal{C}_5^{**} = \mathcal{C}_5^* + \frac{7}{108 \times 25} \Delta_5^*$ with

$$\begin{cases} x(\mathcal{C}_5^{**}) = \frac{40}{108}, \\ a(\mathcal{C}_5^{**}) = \frac{17}{108}, \\ b(\mathcal{C}_5^{**}) = \frac{-4}{108}, \\ y(\mathcal{C}_5^{**}) = \frac{-5}{108}. \end{cases}$$

- For $k = 7$, $g_s(\mathcal{C}_k^*)$ achieves the maximum $\frac{55}{49}$ at $\mathcal{S}_7(5, 5)$ and the second largest value $\frac{54}{49}$ at $\mathcal{S}_7(6, 6)$. By Definition 5.15, all these samples are of class I. Then, an application of Lemma 5.17 yields that

$$h_s(\varepsilon \Delta_7^*) = \begin{cases} -\frac{60}{11}\varepsilon, & s \in \mathcal{S}_7(5, 5), \\ \frac{-98\gamma(s)+48}{11}\varepsilon, & s \in \mathcal{S}_7(6, 6), \end{cases}$$

based on which we infer that the second largest value increases the fastest (with speed $h_s(\varepsilon \Delta_7^*) = \frac{48\varepsilon}{11}$) when $\gamma(s) = 0$. Solving the equation $\frac{55}{49} - \frac{60}{11}\varepsilon = \frac{54}{49} + \frac{48}{11}\varepsilon$, we have $\varepsilon^* = \frac{11}{108 \times 49}$ and obtain $\mathcal{C}_7^{**} = \mathcal{C}_7^* + \frac{11}{108 \times 49} \Delta_7^*$ with

$$\begin{cases} x(\mathcal{C}_7^{**}) = \frac{30}{108}, \\ a(\mathcal{C}_7^{**}) = \frac{13}{108}, \\ b(\mathcal{C}_7^{**}) = \frac{2}{108}, \\ y(\mathcal{C}_7^{**}) = \frac{-3}{108}. \end{cases}$$

- For $k = 8$, $g_s(\mathcal{C}_k^*)$ achieves the maximum $\frac{9}{8}$ at $\mathcal{S}_8(6, 6)$ and the second largest value $\frac{35}{32}$ at $\mathcal{S}_8(5, 5) \cup \mathcal{S}_8(7, 7)$. By Definition 5.15, all these samples are of class I. Then, an application of Lemma 5.17 yields that

$$h_s(\varepsilon \Delta_8^*) = \begin{cases} -\frac{24}{5}\varepsilon, & s \in \mathcal{S}_8(6, 6), \\ \frac{32\gamma(s)-50}{5}\varepsilon, & s \in \mathcal{S}_8(5, 5), \\ \frac{-32\gamma(s)+14}{11}\varepsilon, & s \in \mathcal{S}_8(7, 7), \end{cases}$$

based on which we infer that the second largest value increases the fastest (with speed $h_s(\varepsilon \Delta_8^*) = \frac{110\varepsilon}{5}$) when $s \in \mathcal{S}_8(5, 5, 5)$. Solving the equation $\frac{9}{8} - \frac{24}{5}\varepsilon = \frac{35}{32} + \frac{110}{5}\varepsilon$, we have $\varepsilon^* = \frac{5}{134 \times 32}$ and obtain $\mathcal{C}_8^{**} = \mathcal{C}_8^* + \frac{5}{134 \times 32} \Delta_8^*$ with

$$\begin{cases} x(\mathcal{C}_8^{**}) = \frac{65}{268}, \\ a(\mathcal{C}_8^{**}) = \frac{29}{268}, \\ b(\mathcal{C}_8^{**}) = \frac{-4}{268}, \\ y(\mathcal{C}_8^{**}) = \frac{-5}{268}. \end{cases}$$

- For $k = 9$, $g_s(\mathcal{C}_k^*)$ achieves the maximum $\frac{91}{81}$ at $\mathcal{S}_9(7, 7)$ and the second largest value $\frac{90}{81}$ at $\mathcal{S}_9(5, 5) \cup \mathcal{S}_9(6, 6)$. By Definition 5.15, all these samples are of class I. Then, an application of Lemma 5.17 yields that

$$h_s(\varepsilon \Delta_9^*) = \begin{cases} -\frac{140}{31}\varepsilon, & s \in \mathcal{S}_9(7, 7), \\ \frac{168\gamma(s)-288}{31}\varepsilon, & s \in \mathcal{S}_9(6, 6), \end{cases}$$

based on which, we infer that the second largest value increased fastest (with speed $h_s(\varepsilon \Delta_9^*) = \frac{684\varepsilon}{31}$) when $s \in \mathcal{S}_9(6, 6, 6)$. Solving the equation $\frac{91}{81} - \frac{140}{31}\varepsilon = \frac{90}{81} + \frac{684}{31}\varepsilon$, we have $\varepsilon^* = \frac{31}{824 \times 81}$ and obtain $\mathcal{C}_9^{**} = \mathcal{C}_9^* + \frac{31}{824 \times 81} \Delta_9^*$ with

$$\begin{cases} x(\mathcal{C}_9^{**}) = \frac{176}{824}, \\ a(\mathcal{C}_9^{**}) = \frac{81}{824}, \\ b(\mathcal{C}_9^{**}) = \frac{-10}{824}, \\ y(\mathcal{C}_9^{**}) = \frac{-11}{824}. \end{cases}$$

Step 3: Compute $\mathcal{S}_k^{max}(\mathcal{C}_k^{})$.**

- For $k = 5$, it can be easily verified that $\varepsilon = \frac{7}{108 \times 25}$ is $g_{\mathcal{S}_5}$ -valid. Hence, we compute $g_s(\mathcal{C}_5^{**}) = g_s(\mathcal{C}_5^*) + h_s(\varepsilon \Delta_5^*)$ for all $s \in \mathcal{S}_5$ using Lemma 4.3 and Lemma 5.17. It turns out $g_{\mathcal{S}_5}(\mathcal{C}_5^{**}) = \frac{28}{25} - \frac{24}{2700} = \frac{10}{9}$ is achieved at $\mathcal{S}_5^{max}(\mathcal{C}_5^{**}) = \mathcal{S}_5(3, 3, 3) \cup \mathcal{S}_5(4, 4)$.
- For $k = 7$, it can be easily verified that $\varepsilon = \frac{11}{108 \times 49}$ is not g_s -valid if and only if

$$\begin{aligned} T(s) \in & \left\{ \begin{bmatrix} 2 \\ 1 \end{bmatrix}, \begin{bmatrix} 2 \\ 0 \end{bmatrix}, \begin{bmatrix} 2 \\ 0 \end{bmatrix}, \begin{bmatrix} 2 \\ 0 \end{bmatrix}, \begin{bmatrix} 2 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \end{bmatrix} \right\} \cup \left\{ \begin{bmatrix} 2 \\ 0 \end{bmatrix}, \begin{bmatrix} 2 \\ 0 \end{bmatrix}, \begin{bmatrix} 2 \\ 0 \end{bmatrix}, \begin{bmatrix} 2 \\ 0 \end{bmatrix}, \begin{bmatrix} 2 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right\} \\ & \cup \left\{ \begin{bmatrix} 2 \\ 1 \end{bmatrix}, \begin{bmatrix} 2 \\ 0 \end{bmatrix}, \begin{bmatrix} 2 \\ 0 \end{bmatrix}, \begin{bmatrix} 2 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right\} \cup \left\{ \begin{bmatrix} 2 \\ 0 \end{bmatrix}, \begin{bmatrix} 2 \\ 0 \end{bmatrix}, \begin{bmatrix} 2 \\ 0 \end{bmatrix}, \begin{bmatrix} 2 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 2 \end{bmatrix} \right\} \\ & \cup \left\{ \begin{bmatrix} 2 \\ 1 \end{bmatrix}, \begin{bmatrix} 2 \\ 1 \end{bmatrix}, \begin{bmatrix} 2 \\ 0 \end{bmatrix}, \begin{bmatrix} 2 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \end{bmatrix} \right\} \cup \left\{ \begin{bmatrix} 2 \\ 2 \end{bmatrix}, \begin{bmatrix} 2 \\ 0 \end{bmatrix}, \begin{bmatrix} 2 \\ 0 \end{bmatrix}, \begin{bmatrix} 2 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \end{bmatrix} \right\} \\ & \cup \left\{ \begin{bmatrix} 2 \\ 1 \end{bmatrix}, \begin{bmatrix} 2 \\ 0 \end{bmatrix}, \begin{bmatrix} 2 \\ 0 \end{bmatrix}, \begin{bmatrix} 2 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 2 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \end{bmatrix} \right\}. \end{aligned}$$

It turns out $g_s(\mathcal{C}_7^{**}) \leq \frac{29}{27}$ for all the samples s of the types as above. We compute $g_s(\mathcal{C}_7^{**}) = g_s(\mathcal{C}_7^*) + h_s(\varepsilon \Delta_7^*)$ for all the other samples s using Lemma 4.3 and Lemma 5.17. It turns out $g_{\mathcal{S}_7}(\mathcal{C}_7^{**}) = \frac{55}{49} - \frac{60}{108 \times 49} = \frac{10}{9}$ is achieved at $\mathcal{S}_7^{max}(\mathcal{C}_7^{**}) = \mathcal{S}_7(4, 4, 4) \cup \mathcal{S}_7(5, 5) \cup \mathcal{S}_7(6, 6, 0)$.

- For $k = 8$, it can be easily verified that $\varepsilon = \frac{11}{108 \times 49}$ is not g_s -valid if and only if

$$T(s) \in \left\{ \begin{bmatrix} 2 \\ 0 \end{bmatrix}, \begin{bmatrix} 2 \\ 0 \end{bmatrix}, \begin{bmatrix} 2 \\ 0 \end{bmatrix}, \begin{bmatrix} 2 \\ 0 \end{bmatrix}, \begin{bmatrix} 2 \\ 0 \end{bmatrix}, \begin{bmatrix} 2 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right\} \cup \left\{ \begin{bmatrix} 2 \\ 1 \end{bmatrix}, \begin{bmatrix} 2 \\ 0 \end{bmatrix}, \begin{bmatrix} 2 \\ 0 \end{bmatrix}, \begin{bmatrix} 2 \\ 0 \end{bmatrix}, \begin{bmatrix} 2 \\ 0 \end{bmatrix}, \begin{bmatrix} 2 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \end{bmatrix} \right\}.$$

It turns out $g_s(\mathcal{C}_8^{**}) \leq \frac{71}{67}$ for all the samples s of the types as above. We compute $g_s(\mathcal{C}_8^{**}) = g_s(\mathcal{C}_8^*) + h_s(\varepsilon \Delta_8^*)$ for all the other $s \in \mathcal{S}_8$ using Lemma 4.3 and Lemma 5.17. It turns out $g_{\mathcal{S}_8}(\mathcal{C}_8^{**}) = \frac{9}{8} - \frac{24}{134 \times 32} = \frac{75}{67}$ is achieved at $\mathcal{S}_8^{max}(\mathcal{C}_8^{**}) = \mathcal{S}_8(5, 5, 5) \cup \mathcal{S}_8(6, 6)$.

- For $k = 9$, it can be easily verified that $\varepsilon = \frac{31}{824 \times 81}$ is $g_{\mathcal{S}_9}$ -valid. We compute $g_s(\mathcal{C}_9^{**}) = g_s(\mathcal{C}_9^*) + h_s(\varepsilon \Delta_9^*)$ for all $s \in \mathcal{S}_9$ using Lemma 4.3 and Lemma 5.17. It turns out $g_{\mathcal{S}_9}(\mathcal{C}_9^{**}) = \frac{91}{81} - \frac{140}{824 \times 81} = \frac{231}{206}$ is achieved at $\mathcal{S}_9^{max}(\mathcal{C}_9^{**}) = \mathcal{S}_9(6, 6, 6) \cup \mathcal{S}_9(7, 7)$.

6.2 Optimality of \mathcal{C}_k^{**} for $k = 3, 4, 5, 7, 8, 9$

In this section, we prove that \mathcal{C}_k^{**} obtained in the last section are optimal solutions to $\mathcal{P}_{\mathcal{S}_k}$ for $k = 3, 4, 5, 7, 8, 9$. We first introduce more notations and state some needed lemmas.

Recall that for any sample $s \in \mathcal{S}_k^\circ(\mathcal{C})$, we have $g_s^{(\ell)}(\mathcal{C}) > 0$ for any $\ell \in \text{Ind}_s$, and $g_s^{(\ell)}(\mathcal{C}) < 0$ for any $\ell \notin \text{Ind}_s$. Since any function $g_s^{(\ell)}$, $s \in \mathcal{S}_k^\circ(\mathcal{C})$, is continuous, there exists a neighborhood, denoted by $N(\mathcal{C}, \varepsilon) \subset \mathfrak{S}_k$, of \mathcal{C} such that for all $\mathcal{C}' \in N(\mathcal{C}, \varepsilon)$, all $s \in \mathcal{S}_k(\mathcal{C})$ and all $1 \leq \ell \leq k$,

$$g_s^{(\ell)}(\mathcal{C}') \cdot g_s^{(\ell)}(\mathcal{C}) > 0.$$

For $\mathcal{C}' \in N(\mathcal{C}, \varepsilon)$, we write

$$\Delta := \mathcal{C}' - \mathcal{C} = (\Delta^{(1)}, \Delta^{(2)}, \dots, \Delta^{(k)}), \quad (35)$$

where each $\Delta^{(\ell)} = \left(\delta_{i,j}^{(\ell)} \right)$ is a $k \times k$ matrix such that for all $\ell = 1, 2, \dots, k$,

$$\sum_{i=1}^k \delta_{i,j}^{(\ell)} = \sum_{i=1}^k \delta_{i,j}^{(\ell)} = 0. \quad (36)$$

And moreover, we write

$$h_s(\Delta) := g_s(\mathcal{C}') - g_s(\mathcal{C}). \quad (37)$$

We need the following three lemmas.

Lemma 6.1. *Let $k \geq 3$ and $d \geq 2$. If $\mathcal{S}_k(d, d, d) \subseteq \mathcal{S}_k^\dagger(\mathcal{C})$, then*

$$\sum_{s \in \mathcal{S}_k(d, d, d)} h_s(\Delta) = \left[\sum_{i=1}^k \delta_{i,i}^{(i)} \quad \sum_{i,j:i \neq j} \delta_{j,j}^{(i)} \right] \begin{bmatrix} A \\ B \end{bmatrix}, \quad (38)$$

where $A = \binom{k-1}{d-1}$, $B = \binom{k-2}{d-2} - \binom{k-2}{d-1}$.

Proof. Note that

$$\sum_{s \in \mathcal{S}_k(d, d, d)} h_s(\Delta) = \sum_{\ell=1}^k \sum_{i=1}^k \sum_{j=1}^k h_{i,j}^{(\ell)} \delta_{i,j}^{(\ell)},$$

where the coefficients $h_{i,j}^{(\ell)}$ can be computed as follows. Firstly, note that for any $s \in \mathcal{S}_k(d, d, d)$ and any $(i, j) \in [k] \times [k]$ with $i \neq j$, we have $(i, j) \notin s$, and hence $h_{i,j}^{(\ell)} = 0$. Secondly, for each $(i, i) \in [k] \times [k]$, there are $\binom{k-1}{d-1}$ samples of $\mathcal{S}_k(d, d, d)$ containing (i, i) and hence $h_{i,i}^{(i)} = \binom{k-1}{d-1}$ for all $1 \leq i \leq k$. Thirdly, noticing that for any $j \neq i$, there are $\binom{k-2}{d-2}$ samples containing (i, i) and (j, j) , and there are $\binom{k-2}{d-1}$ samples containing (i, i) but not (j, j) , we have $h_{i,i}^{(j)} = \binom{k-2}{d-2} - \binom{k-2}{d-1}$, which completes the proof. \square

Lemma 6.2. *Let $k \geq 3$ and $d \geq 3$. If $\mathcal{S}_k(d, d, 0) \subseteq \mathcal{S}_k^\dagger(\mathcal{C})$, then*

$$\sum_{s \in \mathcal{S}_k(d, d, 0)} h_s(\Delta) = \left[\sum_{i=1}^k \delta_{i,i}^{(i)} \quad \sum_{i,j:i \neq j} \delta_{j,j}^{(i)} \right] \begin{bmatrix} A' \\ B' \end{bmatrix}, \quad (39)$$

where $A' = (d-1)^{(d-1)} \cdot \left(\binom{k-3}{d-2} - \binom{k-3}{d-3} - 2\binom{k-2}{d-2} \right)$, $B' = (d-1)^{(d-1)} \cdot \left(\binom{k-3}{d-2} - \binom{k-3}{d-3} \right)$.

Proof. Note that

$$\sum_{s \in \mathcal{S}_k(d, d, 0)} h_s(\Delta) = \sum_{\ell=1}^k \sum_{i=1}^k \sum_{j=1}^k h_{i,j}^{(\ell)} \delta_{i,j}^{(\ell)},$$

where the coefficients $h_{i,j}^{(\ell)}$ can be computed as follows. Firstly, note that for any $s \in \mathcal{S}_k(d, d, 0)$, we have $(i, i) \notin s$, and hence $h_{i,i}^{(\ell)} = 0$ for all $1 \leq i, \ell \leq k$. Secondly, for each $i \neq j$, there are $(d-1)^{(d-1)} \cdot \binom{k-2}{d-2}$ samples from $\mathcal{S}_k(d, d, 0)$ containing (i, j) and hence $h_{i,j}^{(i)} = h_{i,j}^{(j)} = (d-1)^{(d-1)} \cdot \binom{k-2}{d-2}$. Thirdly, noticing that for any distinct i, j, ℓ , there are $(d-1)^{(d-1)} \cdot \binom{k-3}{d-3}$ samples s such that $\ell \in \text{Ind}_s$ and $(i, j) \in s$, and there are $(d-1)^{(d-1)} \cdot \binom{k-3}{d-2}$ samples s such that $\ell \notin \text{Ind}_s$ and $(i, j) \in s$, we have $h_{i,j}^{(\ell)} = (d-1)^{(d-1)} \cdot \left(\binom{k-3}{d-3} - \binom{k-3}{d-2} \right)$. Finally, the desired result follows from (36). \square

Lemma 6.3. *If $\mathcal{S}_3(2, 2, 0) \subseteq \mathcal{S}_3^\dagger(\mathcal{C})$, then*

$$\sum_{s \in \mathcal{S}_3(2,2,0)} h_s(\Delta) = \left[\sum_{i=1}^3 \delta_{i,i}^{(i)} \quad \sum_{i,j:i \neq j} \delta_{j,j}^{(i)} \right] \begin{bmatrix} -3 \\ 1 \end{bmatrix}. \quad (40)$$

Proof. For the 3 samples in $\mathcal{S}_3(2, 2, 0)$, we have

$$\begin{aligned} h_{\{(2,1),(1,2)\}}(\Delta) &= (\delta_{2,1}^{(1)} + \delta_{1,2}^{(1)}) + (\delta_{2,1}^{(2)} + \delta_{1,2}^{(2)}) - (\delta_{2,1}^{(3)} + \delta_{1,2}^{(3)}), \\ h_{\{(3,2),(2,3)\}}(\Delta) &= (\delta_{3,2}^{(2)} + \delta_{2,3}^{(2)}) + (\delta_{3,2}^{(3)} + \delta_{2,3}^{(3)}) - (\delta_{3,2}^{(1)} + \delta_{2,3}^{(1)}), \\ h_{\{(3,1),(1,3)\}}(\Delta) &= (\delta_{3,1}^{(1)} + \delta_{1,3}^{(1)}) + (\delta_{3,1}^{(3)} + \delta_{1,3}^{(3)}) - (\delta_{3,1}^{(2)} + \delta_{1,3}^{(2)}). \end{aligned}$$

Hence,

$$\begin{aligned} \sum_{s \in \mathcal{S}_3(2,2,0)} h_s(\Delta) &= \sum_{\ell=1}^3 \sum_{i \neq j} \delta_{i,j}^{(\ell)} - 2 \sum_{\text{distinct } i,j,\ell} \delta_{i,j}^{(\ell)} \\ &= \sum_{i=1}^3 -3\delta_{i,i}^{(i)} + \sum_{i,j:i \neq j} \delta_{j,j}^{(i)}, \end{aligned} \quad (41)$$

which complete the proof. \square

The following lemma gives a sufficient condition for the local optimality of an arbitrary $\mathcal{C} \in \mathfrak{S}_k$.

Lemma 6.4. *Let $\mathcal{C} \in \mathfrak{S}_k$. If there exists a subset $\mathcal{S}_k^\circ(\mathcal{C}) \subseteq \mathcal{S}_k^\dagger(\mathcal{C})$ and a neighborhood $N(\mathcal{C}, \varepsilon) \subset \mathfrak{S}_k$ of \mathcal{C} and a set of positive reals $\{k_s \mid s \in \mathcal{S}_k^\circ(\mathcal{C})\}$ such that for all $\mathcal{C}' \in N(\mathcal{C}, \varepsilon)$,*

$$\sum_{s \in \mathcal{S}_k^\circ(\mathcal{C})} k_s \cdot (g_s(\mathcal{C}') - g_s(\mathcal{C})) = 0. \quad (42)$$

Then, \mathcal{C} is a local optimal point for $\mathcal{P}_{\mathcal{S}_k}$.

Proof. Suppose, by way of contradiction, that \mathcal{C} is not a local optimal point, i.e., there exists a neighborhood $N(\mathcal{C}, \varepsilon)$ of \mathcal{C} and $\mathcal{C}' \in N(\mathcal{C}, \varepsilon)$ such that $g_{\mathcal{S}_k}(\mathcal{C}') < g_{\mathcal{S}_k}(\mathcal{C})$. Then, for all $s \in \mathcal{S}_k(\mathcal{C})$,

$$g_s(\mathcal{C}') \leq \max_{s \in \mathcal{S}_k} \{g_s(\mathcal{C}')\} = g_{\mathcal{S}_k}(\mathcal{C}') < g_{\mathcal{S}_k}(\mathcal{C}) = \max_{s \in \mathcal{S}_k} \{g_s(\mathcal{C})\} = g_s(\mathcal{C}),$$

which contradicts (42). Hence, \mathcal{C} is an local optimal point. \square

From now on, we denote the vector $\begin{bmatrix} A \\ B \end{bmatrix}$ in (38) by $\mathcal{H}_k(d, d, d)$, $\frac{1}{(d-1)^{(d-1)}} \begin{bmatrix} A' \\ B' \end{bmatrix}$ by $\mathcal{H}_k(d, d, 0)$ where $\begin{bmatrix} A' \\ B' \end{bmatrix}$ is obtained in (39) and $\mathcal{H}_3(2, 2, 0) = \begin{bmatrix} -3 \\ 1 \end{bmatrix}$ by (41). We are then ready to give the main result of this section.

Theorem 6.5. *\mathcal{C}_k^{**} is an optimal point for $\mathcal{P}_{\mathcal{S}_k}$ for $k = 3, 4, 5, 7, 8, 9$.*

Proof. By Lemma 2.2, it suffices to prove that \mathcal{C}_k^{**} is a local optimal point. To this end, by Lemma 6.4, we only need to find a neighborhood of \mathcal{C}_k^{**} , a subset $\mathcal{S}_k^\circ(\mathcal{C}_k^{**})$ of $\mathcal{S}_k^\dagger(\mathcal{C}_k^{**})$ and a set of positives reals satisfying (42). In the following, we take $N(\mathcal{C}_k^{**}, \varepsilon)$ as the neighborhood of \mathcal{C}_k^{**} for each $k = 3, 4, 5, 7, 8, 9$.

- For the case $k = 3$, let $\mathcal{S}_3^\circ(\mathcal{C}_3^{**}) = \mathcal{S}_3(2, 2, 0) \cup \mathcal{S}_3(2, 2, 2) \cup \mathcal{S}_3(3, 3, 0)$. It can be verified that $\mathcal{S}_3^\circ(\mathcal{C}_3^{**}) \subseteq \mathcal{S}_3^\dagger(\mathcal{C}_3^{**})$. Then, by Lemmas 6.3, 6.1 and 6.2, we infer that (40), (38) and (39) hold with $\mathcal{H}_3(2, 2, 0) = \begin{bmatrix} -3 \\ 1 \end{bmatrix}$, $\mathcal{H}_3(2, 2, 2) = \begin{bmatrix} 2 \\ 0 \end{bmatrix}$ and $\mathcal{H}_3(3, 3, 0) = \begin{bmatrix} -1 \\ -1 \end{bmatrix}$, respectively. Since $\mathcal{H}_3(2, 2, 0) + 2\mathcal{H}_3(2, 2, 2) + \mathcal{H}_3(3, 3, 0) = 0$, an application of Lemma 6.4 yields that \mathcal{C}_3^{**} is an optimal point for $\mathcal{P}_{\mathcal{S}_3}$.
- For the case $k = 4$, let $\mathcal{S}_4^\circ(\mathcal{C}_4^{**}) = \mathcal{S}_4(2, 2, 2) \cup \mathcal{S}_4(3, 3, 0) \cup \mathcal{S}_4(3, 3, 3)$, which can be verified to be a subset of $\mathcal{S}_4^\dagger(\mathcal{C}_4^{**})$. Then, as in previous case, the desired optimality of \mathcal{C}_4^{**} then follows from Lemmas 6.1, 6.2 and 6.4 and the easily verifiable fact that $2\mathcal{H}_4(2, 2, 2) + 3\mathcal{H}_4(3, 3, 0) + 2\mathcal{H}_4(3, 3, 3) = 0$, where $\mathcal{H}_4(2, 2, 2) = \begin{bmatrix} 3 \\ -1 \end{bmatrix}$, $\mathcal{H}_4(3, 3, 0) = \begin{bmatrix} -4 \\ 0 \end{bmatrix}$ and $\mathcal{H}_4(3, 3, 3) = \begin{bmatrix} 3 \\ 1 \end{bmatrix}$.
- For the case $k = 5$, let $\mathcal{S}_5^\circ(\mathcal{C}_5^{**}) = \mathcal{S}_5(3, 3, 3) \cup \mathcal{S}_5(4, 4, 0) \cup \mathcal{S}_5(4, 4, 4) \subseteq \mathcal{S}_5^\dagger(\mathcal{C}_5^{**})$. Then, the desired optimality of \mathcal{C}_5^{**} then follows from Lemmas 6.1, 6.2 and 6.4 and the easily verifiable fact that $\mathcal{H}_5(3, 3, 3) + 2\mathcal{H}_5(4, 4, 0) + \mathcal{H}_5(4, 4, 4) = 0$.
- For the case $k = 7$, let $\mathcal{S}_7^\circ(\mathcal{C}_7^{**}) = \mathcal{S}_7(4, 4, 4) \cup \mathcal{S}_7(5, 5, 0) \cup \mathcal{S}_7(5, 5, 5) \subseteq \mathcal{S}_7^\dagger(\mathcal{C}_7^{**})$. Then, the desired optimality of \mathcal{C}_7^{**} then follows from the fact that $3\mathcal{H}_7(4, 4, 4) + 5\mathcal{H}_7(5, 5, 0) + 2\mathcal{H}_7(5, 5, 5) = 0$.
- For the case $k = 8$, let $\mathcal{S}_8^\circ(\mathcal{C}_8^{**}) = \mathcal{S}_8(5, 5, 5) \cup \mathcal{S}_8(6, 6, 0) \cup \mathcal{S}_8(6, 6, 6) \subseteq \mathcal{S}_8^\dagger(\mathcal{C}_8^{**})$. Then, the desired optimality then follows from the fact that $12\mathcal{H}_8(5, 5, 5) + 21\mathcal{H}_8(6, 6, 0) + 5\mathcal{H}_8(6, 6, 6) = 0$.
- For the case $k = 9$, let $\mathcal{S}_9^\circ(\mathcal{C}_9^{**}) = \mathcal{S}_9(6, 6, 6) \cup \mathcal{S}_9(7, 7, 0) \cup \mathcal{S}_9(7, 7, 7) \subseteq \mathcal{S}_9^\dagger(\mathcal{C}_9^{**})$. Then, the desired optimality then follows the fact that $15\mathcal{H}_9(6, 6, 6) + 28\mathcal{H}_9(7, 7, 0) + 3\mathcal{H}_9(7, 7, 7) = 0$.

□

6.3 The Uniqueness of Optimal Solutions for $k = 3, 4, 5, 7, 8, 9$

We are concerned with the uniqueness of the optimal solutions to $\mathcal{P}_{\mathcal{S}_k}$. Note that the case of $k = 1$ is trivial, and it is known from the proof of Theorem 3 of [20] that \mathcal{C}_2^* is the unique optimal point for $\mathcal{P}_{\mathcal{S}_2}$. In this section, we will show that the optimal solutions to $\mathcal{P}_{\mathcal{S}_k}$ are unique for $k = 3, 4, \dots, 9$, which however ceases to hold true for $k = 10$.

We first need the following lemma, which strengthens Lemma 6.4.

Lemma 6.6. *Let $\mathcal{C} \in \mathfrak{S}_k$. If there exists a subset $\mathcal{S}_k^*(\mathcal{C}) \subseteq \mathcal{S}_k^\dagger(\mathcal{C})$ and a neighborhood $N(\mathcal{C}, \varepsilon) \subset \mathfrak{S}_k$ of \mathcal{C} and a set of positive reals $\{k_s \mid s \in \mathcal{S}_k^*(\mathcal{C})\}$ such that (1) For all $\mathcal{C}' \in N(\mathcal{C}, \varepsilon)$, $\sum_{s \in \mathcal{S}_k^*(\mathcal{C})} k_s \cdot h_s(\Delta) = 0$; (2) If for all $s \in \mathcal{S}_k^*(\mathcal{C})$, $h_s(\Delta) = 0$ then $\Delta = 0$, where, as before, $\Delta = \mathcal{C}' - \mathcal{C}$ and $h_s(\Delta) = g_s(\mathcal{C}') - g_s(\mathcal{C})$. Then, \mathcal{C} is the unique local optimal point for $\mathcal{P}_{\mathfrak{S}_k}$.*

Proof. Suppose, by way of contradiction, that there exists another optimal point $\mathcal{C}' \in N(\mathcal{C}, \varepsilon)$ such that $\mathcal{C}' - \mathcal{C} = \Delta \neq 0$. By Condition (1), we know that for all $s \in \mathcal{S}_k^*(\mathcal{C})$, $g_s(\mathcal{C}') = g_s(\mathcal{C})$, i.e., $h_s(\Delta) = 0$ (Since otherwise there exist $s_0, s_1 \in \mathcal{S}_k^*(\mathcal{C})$ such that $g_{s_0}(\mathcal{C}') - g_{s_0}(\mathcal{C}) > 0$ and $g_{s_1}(\mathcal{C}') - g_{s_1}(\mathcal{C}) < 0$, which contradicts the optimality of \mathcal{C}). Hence, by Condition (2), we have $\Delta = 0$, which contradicts the assumption that $\Delta \neq 0$ and thereby the result follows. \square

In the following, we set

$$\mathcal{S}_3^*(\mathcal{C}_3^{**}) := \mathcal{S}_3(2, 2) \cup \mathcal{S}_3(3, 3, 0).$$

And it can be easily verified that $\mathcal{S}_3^*(\mathcal{C}_3^{**}) \subseteq \mathcal{S}_3^\dagger(\mathcal{C}_3^{**})$. The following lemma can be used to establish the uniqueness of \mathcal{C}_3^{**} for $\mathcal{P}_{\mathfrak{S}_3}$.

Lemma 6.7. *There exist a neighborhood $N(\mathcal{C}_3^{**}, \varepsilon) \subset \mathfrak{S}_3$ of \mathcal{C}_3^{**} and a set of positive reals $\{k_s \mid s \in \mathcal{S}_3^*(\mathcal{C}_3^{**})\}$ such that for all $\mathcal{C}' \in N(\mathcal{C}_3^{**}, \varepsilon)$*

$$\sum_{s \in \mathcal{S}_3^*(\mathcal{C}_3^{**})} k_s \cdot (g_s(\mathcal{C}') - g_s(\mathcal{C}_3^{**})) = 0. \quad (43)$$

Proof. From the proof of Theorem 6.5, we have that for all $\mathcal{C}' \in N(\mathcal{C}_3^{**}, \varepsilon)$,

$$\sum_{s \in \mathcal{S}_3(2, 2, 0)} h_s(\Delta) + 2 \sum_{s \in \mathcal{S}_3(2, 2, 2)} h_s(\Delta) + \sum_{s \in \mathcal{S}_3(3, 3, 0)} h_s(\Delta) = 0,$$

where, as before, $\Delta = \mathcal{C}' - \mathcal{C}_3^{**}$. Since

$$\begin{aligned} h_{\{(1,1),(1,2)\}}(\Delta) + h_{\{(2,1),(2,2)\}}(\Delta) &= h_{\{(1,1),(2,2)\}}(\Delta) + h_{\{(2,1),(1,2)\}}(\Delta), \\ h_{\{(1,1),(1,3)\}}(\Delta) + h_{\{(3,1),(3,3)\}}(\Delta) &= h_{\{(1,1),(3,3)\}}(\Delta) + h_{\{(3,1),(1,3)\}}(\Delta), \\ h_{\{(2,2),(2,3)\}}(\Delta) + h_{\{(3,2),(3,3)\}}(\Delta) &= h_{\{(2,2),(3,3)\}}(\Delta) + h_{\{(3,2),(2,3)\}}(\Delta), \end{aligned}$$

we have

$$\sum_{s \in \mathcal{S}_3(2, 2, 1)} h_s(\Delta) = \sum_{s \in \mathcal{S}_3(2, 2, 0)} h_s(\Delta) + \sum_{s \in \mathcal{S}_3(2, 2, 2)} h_s(\Delta). \quad (44)$$

Hence, we have

$$\frac{1}{2} \sum_{s \in \mathcal{S}_3(2, 2, 0)} h_s(\Delta) + \frac{1}{2} \sum_{s \in \mathcal{S}_3(2, 2, 1)} h_s(\Delta) + \frac{3}{2} \sum_{s \in \mathcal{S}_3(2, 2, 2)} h_s(\Delta) + \sum_{s \in \mathcal{S}_3(3, 3, 0)} h_s(\Delta) = 0.$$

The proof is then complete. \square

In the following, we set

$$\mathcal{S}_4^*(\mathcal{C}_4^{**}) := \mathcal{S}_4(3, 3, 0) \cup \mathcal{S}_4(3, 3, 1) \cup \mathcal{S}_4(4, 4, 0).$$

It can be easily verified that $\mathcal{S}_4^*(\mathcal{C}_4^{**}) \subseteq \mathcal{S}_4^\dagger(\mathcal{C}_4^{**})$. The following lemma, whose proof has been postponed to Appendix C, can be used to establish the uniqueness of \mathcal{C}_4^{**} for $\mathcal{P}_{\mathcal{S}_4}$.

Lemma 6.8. *There exists a neighborhood $N(\mathcal{C}_4^{**}, \varepsilon) \subset \mathfrak{S}_4$ of \mathcal{C}_4^{**} and a set of positive reals $\{k_s \mid s \in \mathcal{S}_4^*(\mathcal{C}_4^{**})\}$ such that for all $\mathcal{C}' \in N(\mathcal{C}_4^{**}, \varepsilon)$,*

$$\sum_{s \in \mathcal{S}_4^*(\mathcal{C}_4^{**})} k_s \cdot (g_s(\mathcal{C}') - g_s(\mathcal{C}_4^{**})) = 0. \quad (45)$$

We are now ready to give the main result of this section.

Theorem 6.9. \mathcal{C}_k^{**} is the unique optimal point for $\mathcal{P}_{\mathcal{S}_k}$ for $k = 3, 4, \dots, 9$.

Proof. ■ We first deal with the case $k = 3$. By Lemma 6.6 and then Lemma 6.7, we only need to prove that the equation

$$h_s(\Delta) = 0 \quad \text{for all } s \in \mathcal{S}_3(\mathcal{C}_3^{**}) \quad (46)$$

has the unique solution $\Delta = 0$.

Suppose $\Delta = \left((\delta_{i,j}^{(1)}), (\delta_{i,j}^{(2)}), (\delta_{i,j}^{(3)}) \right)$ is a solution of (46). We first prove that for all $1 \leq i, j \leq 3$,

$$\pi_{i,j} := \sum_{\ell=1}^3 \delta_{i,j}^{(\ell)} = 0. \quad (47)$$

By (36), we have $\pi_{1,1} + \pi_{1,2} + \pi_{1,3} = 0$ and then by $h_{\{(2,1),(1,2),(1,3)\}}(\Delta) = \pi_{2,1} + \pi_{1,2} + \pi_{1,3} = 0$, we have $\pi_{1,1} = \pi_{2,1}$. Similarly, we have $\pi_{1,1} = \pi_{2,1} = \pi_{3,1}$. By (36), we have $\pi_{1,1} + \pi_{2,1} + \pi_{3,1} = 0$, and hence, $\pi_{1,1} = \pi_{2,1} = \pi_{3,1} = 0$. Further, in the same way, we have $\pi_{1,2} = \pi_{2,2} = \pi_{3,2} = 0$ and finally we obtain (47).

By $h_{\{(1,1),(1,2)\}}(\Delta) = 0$ and (36), we have $\delta_{1,3}^{(3)} = \delta_{1,3}^{(1)} + \delta_{1,3}^{(2)}$. Hence, $0 = \pi_{1,3} = 2\delta_{1,3}^{(3)}$. Similarly, we can have $\delta_{1,3}^{(3)} = \delta_{2,3}^{(3)} = 0$. Hence, by (36), we further have $\delta_{3,3}^{(3)} = 0$. In a similar fashion, we finally have

$$\delta_{i,j}^{(j)} = 0, \quad 1 \leq i, j \leq 3. \quad (48)$$

Letting $\delta_{1,2}^{(1)} = a$, $\delta_{2,2}^{(1)} = b$, $\delta_{3,2}^{(1)} = c$ and using Equations (36), (47) and (48), we have

$$\Delta = \left(\left(\begin{pmatrix} 0 & a & -a \\ 0 & b & -b \\ 0 & c & -c \end{pmatrix}, \begin{pmatrix} -a & 0 & a \\ -b & 0 & b \\ -c & 0 & c \end{pmatrix}, \begin{pmatrix} a & -a & 0 \\ b & -b & 0 \\ c & -c & 0 \end{pmatrix} \right).$$

By $h_{\{(1,1),(2,2)\}}(\Delta) = 0$, we have $b - a = a - b$, i.e., $a = b$. By $h_{\{(1,1),(3,3)\}}(\Delta) = 0$, we have $c - a = a - c$, i.e., $a = c$. Hence $a = b = c$ and then by Equation (36), we have $a = b = c = 0$, which means $\Delta = 0$. The proof is then complete.

■ We now deal with the case $k = 4$. By Lemma 6.6 and then Lemma 6.8, we only need to prove that the equation

$$h_s(\Delta) = 0 \quad \text{for all } s \in \mathcal{S}_4^*(\mathcal{C}_4^{**}) \quad (49)$$

has the unique solution $\Delta = 0$.

Suppose $\Delta = \left((\delta_{i,j}^{(1)}), (\delta_{i,j}^{(2)}), (\delta_{i,j}^{(3)}), (\delta_{i,j}^{(4)}) \right)$ is a solution of (49). We first prove that for all $i, j = 1, 2, 3, 4$,

$$\pi_{i,j} := \sum_{\ell=1}^4 \delta_{i,j}^{(\ell)} = 0. \quad (50)$$

By (36), we have $\pi_{1,1} + \pi_{1,2} + \pi_{1,3} + \pi_{1,4} = 0$ and then by $h_{\{(2,1),(1,2),(1,3),(1,4)\}}(\Delta) = \pi_{2,1} + \pi_{1,2} + \pi_{1,3} + \pi_{1,4} = 0$, we have $\pi_{1,1} = \pi_{2,1}$. Similarly, we have $\pi_{1,1} = \pi_{2,1} = \pi_{3,1} = \pi_{4,1}$. By (36), we have $\pi_{1,1} + \pi_{2,1} + \pi_{3,1} + \pi_{4,1} = 0$, and hence, $\pi_{1,1} = \pi_{2,1} = \pi_{3,1} = \pi_{4,1} = 0$. Further, in the same way, we have $\pi_{1,2} = \pi_{2,2} = \pi_{3,2} = \pi_{4,2} = 0$ and finally, we can obtain (50).

By $h_{\{(1,1),(1,2),(1,3)\}}(\Delta) = 0$ and (36), we have $\delta_{1,4}^{(4)} = \delta_{1,4}^{(1)} + \delta_{1,4}^{(2)} + \delta_{1,4}^{(3)}$. Hence, $0 = \pi_{1,4} = 2\delta_{1,4}^{(4)}$. Similarly, we can have $\delta_{2,4}^{(4)} = \delta_{3,4}^{(4)} = 0$. Hence, by (36), we further have $\delta_{4,4}^{(4)} = 0$. Similarly, we can have

$$\delta_{i,j}^{(j)} = 0, \quad 1 \leq i, j \leq 4. \quad (51)$$

Since $h_{\{(1,1),(1,2),(1,3)\}}(\Delta) = h_{\{(2,1),(1,2),(1,3)\}}(\Delta) = 0$, we have $\delta_{1,1}^{(1)} + \delta_{1,1}^{(2)} + \delta_{1,1}^{(3)} - \delta_{1,1}^{(4)} = \delta_{2,1}^{(1)} + \delta_{2,1}^{(2)} + \delta_{2,1}^{(3)} - \delta_{2,1}^{(4)}$, and furthermore, by (50), $-2\delta_{1,1}^{(4)} = -2\delta_{2,1}^{(4)}$. Similarly, by $h_{\{(1,1),(1,2),(1,3)\}}(\Delta) = h_{\{(3,1),(1,2),(1,3)\}}(\Delta) = 0$, we have

$$\delta_{1,1}^{(4)} = \delta_{2,1}^{(4)} = \delta_{3,1}^{(4)}.$$

Since $h_{\{(1,1),(1,2),(1,4)\}}(\Delta) = h_{\{(2,1),(1,2),(1,4)\}}(\Delta) = 0$, we have $\delta_{1,1}^{(1)} + \delta_{1,1}^{(2)} + \delta_{1,1}^{(4)} - \delta_{1,1}^{(3)} = \delta_{2,1}^{(1)} + \delta_{2,1}^{(2)} + \delta_{2,1}^{(4)} - \delta_{2,1}^{(3)}$, and furthermore, by (50), $-2\delta_{1,1}^{(3)} = -2\delta_{2,1}^{(3)}$. Similarly, by $h_{\{(1,1),(1,2),(1,4)\}}(\Delta) = h_{\{(4,1),(1,2),(1,4)\}}(\Delta) = 0$, we can have

$$\delta_{1,1}^{(3)} = \delta_{2,1}^{(3)} = \delta_{4,1}^{(3)}.$$

Since $h_{\{(1,1),(1,3),(1,4)\}}(\Delta) = h_{\{(3,1),(1,3),(1,4)\}}(\Delta) = 0$, we have $\delta_{1,1}^{(1)} + \delta_{1,1}^{(3)} + \delta_{1,1}^{(4)} - \delta_{1,1}^{(2)} = \delta_{3,1}^{(1)} + \delta_{3,1}^{(3)} + \delta_{3,1}^{(4)} - \delta_{3,1}^{(2)}$, i.e., $-2\delta_{1,1}^{(2)} = -2\delta_{3,1}^{(2)}$ by (50). Similarly, by $h_{\{(1,1),(1,3),(1,4)\}}(\Delta) = h_{\{(4,1),(1,3),(1,4)\}}(\Delta) = 0$, we can have

$$\delta_{1,1}^{(2)} = \delta_{3,1}^{(2)} = \delta_{4,1}^{(2)}.$$

Now, note that $\pi_{1,1} = \delta_{1,1}^{(1)} + \delta_{1,1}^{(2)} + \delta_{1,1}^{(3)} + \delta_{1,1}^{(4)} = \delta_{2,1}^{(1)} + \delta_{2,1}^{(2)} + \delta_{2,1}^{(3)} + \delta_{2,1}^{(4)} = \pi_{2,1}$, by (51) and the above discussions, we have $\delta_{2,1}^{(2)} = \delta_{1,1}^{(2)}$. Similarly, by $\pi_{1,1} = \pi_{3,1}$, we have $\delta_{3,1}^{(3)} = \delta_{1,1}^{(3)}$ and by $\pi_{1,1} = \pi_{4,1}$, we have $\delta_{4,1}^{(4)} = \delta_{1,1}^{(4)}$. Hence, by (36), we deduce that for $i = 1, 2, 3, 4$,

$$\delta_{1,1}^{(i)} = \delta_{2,1}^{(i)} = \delta_{3,1}^{(i)} = \delta_{4,1}^{(i)} = 0.$$

Similarly, one can have that for $i = 1, 2, 3, 4, j = 2, 3, 4$,

$$\delta_{1,j}^{(i)} = \delta_{2,j}^{(i)} = \delta_{3,j}^{(i)} = \delta_{4,j}^{(i)} = 0.$$

Collecting all the results above, we conclude that $\Delta = 0$, as desired.

■ The uniqueness of the optimal solutions for $k = 5, 6, \dots, 9$ follows from a more complex yet completely parallel argument as for $k = 3, 4$, and therefore we omit the details. \square

Theorem 6.10. *There are at least two optimal points for $\mathcal{P}_{\mathcal{S}_{10}}$.*

Proof. It suffices to find an optimal point for $\mathcal{P}_{\mathcal{S}_{10}}$ that is different from \mathcal{C}_{10}^{**} . To this end, consider the system (24) and replace “<” and “>” by “ \leq ” and “ \geq ”, respectively. Then, we have

$$\begin{cases} \bar{a} + \bar{b} \leq 0, \\ \bar{a} + 2\bar{b} \leq 0, \\ \bar{a} + 2\bar{b} \geq 0, \\ 7\bar{a} + 20\bar{b} \geq 0. \end{cases} \quad (52)$$

Note that the above system has solution

$$\begin{cases} \bar{b} \geq 0, \\ \bar{a} + 2\bar{b} = 0. \end{cases} \quad (53)$$

Now choosing $\delta > 0$ small enough and setting $b = \delta$, we obtain an optimal point $\mathcal{C} = \mathcal{C}_{10}^* + \Delta$ different from \mathcal{C}_{10}^{**} with $\bar{a}(\Delta) = -2\delta$, $\bar{b}(\Delta) = \delta$, $\bar{y}(\Delta) = -6\delta$, $\bar{x}(\Delta) = 1 + 18\delta$. \square

6.4 Routing Rate

By Theorem 2.5, the optimal solution \mathcal{C}_k^{**} , $k = 3, 4, \dots, 10$, gives an explicit construction of multi-flows for the corresponding k -pair strongly reachable network. More precisely, translating the results in this section, we have constructed multi-flows of rate $(\frac{11}{12}, \dots, \frac{11}{12})$ for $k = 3, 4$, rate $(\frac{9}{10}, \dots, \frac{9}{10})$ for $k = 5, 6, 7$, rate $(\frac{67}{75}, \dots, \frac{67}{75})$ for $k = 8$, rate $(\frac{206}{231}, \dots, \frac{206}{231})$ for $k = 9$, rate $(\frac{25}{28}, \dots, \frac{25}{28})$ for $k = 10$, each of which further gives a lower bound on the corresponding $\mathbf{R}_r(\mathcal{N})$. To the best of our knowledge, the aforementioned rates are the largest to date.

7 Concluding Remarks

We attack the Langberg-Médard multiple unicast conjecture via an optimization approach. For a closely related optimization problem $\mathcal{P}_{\mathcal{S}_k}$ with optimal value $\mathcal{O}_{\mathcal{S}_k}$, we analyze the asymptotics of $\{\mathcal{O}_{\mathcal{S}_k}\}$ and explicitly solve $\mathcal{P}_{\mathcal{S}_k}$ for $k = 1, 2, \dots, 10$. More precisely, we prove that $\lim_{k \rightarrow \infty} \mathcal{O}_{\mathcal{S}_k} = 9/8$, and establish the first 10 terms of $\{\mathcal{O}_{\mathcal{S}_k}\}$ $1, 1, \frac{12}{11}, \frac{12}{11}, \frac{10}{9}, \frac{10}{9}, \frac{10}{9}, \frac{75}{67}, \frac{231}{206}, \frac{28}{25}$, which give the largest feasible routing rate to date for the corresponding strongly reachable networks.

For any $k \neq 1, 2, 6, 10$, there exists a perturbation promising to give better solutions than \mathcal{C}_k^* , a sequence of asymptotically optimal solutions to $\mathcal{P}_{\mathcal{S}_k}$, and moreover, a delicate perturbation analysis in Sections 5 and 6 gives the exact optimal solutions for $k \leq 10$. Nevertheless, it remains to be seen whether the perturbation approach can be used to solve $\mathcal{P}_{\mathcal{S}_k}$ for all k . The major hurdle for the case of larger k is the drastically increasing complexity needed for the analysis, which is already prohibitive for $k = 11$. Here we remark that the optimization problem appears to be “trickier” than previously thought. For a quick example,

one would be tempted to think that the sequence $\{\mathcal{O}_{\mathcal{S}_k}\}$ should be monotonically increasing. This, however, is not true, since our results actually indicate that $\mathcal{O}_{\mathcal{S}_9} > \mathcal{O}_{\mathcal{S}_{10}}$.

Appendices

A Proof of Lemma 5.8

By the definition of \mathcal{C}_k^* , it can be readily verified that for any 3-sample s and any $1 \leq \ell \leq 3$,

$$g_s^{(\ell)}(\mathcal{C}_3^*) = \frac{3}{9}m_{Ind_s}(\ell) - \frac{1}{9}\alpha(s). \quad (54)$$

And note that for all $s \in \mathcal{S}_3$, $0 \leq m_{Ind_s}(\ell) \leq 4$ and

$$h_s^{(\ell)}(\varepsilon \Delta_k^*) = \varepsilon \cdot \sum_{(i,j) \in s} \delta_{i,j}^{(\ell)},$$

where $\delta_{i,j}^{(\ell)}$ is defined as in $\Delta_k^* = ((\delta_{i,j}^{(1)}), (\delta_{i,j}^{(2)}), \dots, (\delta_{i,j}^{(k)}))$.

By Definition 5.6, any $\varepsilon > 0$ is $g_s^{(1)}$ -valid since $g_s^{(1)}(\mathcal{C}_3^*) = 0$. Recall from Example 5.5 that $\bar{x}(\Delta_3^*) = -2$, $\bar{y}(\Delta_3^*) = -5$, $\bar{a}(\Delta_3^*) = 1$ and $\bar{b}(\Delta_3^*) = 4$. Then, by definition, it can be easily verified that

$$\{g_s^{(\ell)}(\mathcal{C}_3^*) \mid s \in \mathcal{S}_3, 1 \leq \ell \leq 3\} \subseteq \left\{ \frac{-2}{9}, \frac{-1}{9}, 0, \frac{1}{9}, \dots, \frac{8}{9}, 1 \right\}.$$

We now deal with the following cases:

- If $g_s^{(\ell)}(\mathcal{C}_3^*) = \frac{-2}{9}$, which implies $m_{Ind_s}(\ell) = 0$ and $\alpha(s) = 2$ (see Equation (54)), then ε is $g_s^{(\ell)}$ -valid if and only if

$$\begin{cases} 2\bar{y}(\Delta_3^*)\varepsilon \leq \frac{2}{9}, \\ 2\bar{b}(\Delta_3^*)\varepsilon \leq \frac{2}{9}, \\ (\bar{y}(\Delta_3^*) + \bar{b}(\Delta_3^*))\varepsilon \leq \frac{2}{9}. \end{cases}$$

Recalling from Example 5.5 that $\bar{x}(\Delta_3^*) = -2$, $\bar{y}(\Delta_3^*) = -5$, $\bar{a}(\Delta_3^*) = 1$ and $\bar{b}(\Delta_3^*) = 4$, we deduce that for this case $\varepsilon > 0$ is $g_s^{(\ell)}$ -valid if and only if $\varepsilon \leq \frac{1}{36}$.

- If $g_s^{(\ell)}(\mathcal{C}_3^*) = \frac{-1}{9}$, which implies $m_{Ind_s}(\ell) = 0$ and $\alpha(s) = 1$, then ε is $g_s^{(\ell)}$ -valid if and only if

$$\begin{cases} \bar{y}(\Delta_3^*)\varepsilon \leq \frac{1}{9}, \\ \bar{b}(\Delta_3^*)\varepsilon \leq \frac{1}{9}. \end{cases}$$

Similarly, we deduce that for this case $\varepsilon > 0$ is $g_s^{(\ell)}$ -valid if and only if $\varepsilon \leq \frac{1}{36}$.

- If $g_s^{(\ell)}(\mathcal{C}_3^*) = 0$, then, by definition, any $\varepsilon > 0$ is $g_s^{(\ell)}$ -valid.

- If $g_s^{(\ell)}(\mathcal{C}_3^*) = \frac{1}{9}$, which implies $m_{Ind_s}(\ell) = 1$ and $\alpha(s) = 2$, then ε is $g_s^{(\ell)}$ -valid if and only if

$$\begin{cases} -(\bar{a}(\Delta_3^*) + \bar{y}(\Delta_3^*))\varepsilon \leq \frac{1}{9}, \\ -(\bar{a}(\Delta_3^*) + \bar{b}(\Delta_3^*))\varepsilon \leq \frac{1}{9}. \end{cases}$$

Straightforward computations yield that that, for this case, $\varepsilon > 0$ is $g_s^{(\ell)}$ -valid if and only if $\varepsilon \leq \frac{1}{36}$.

- If $g_s^{(\ell)}(\mathcal{C}_3^*) = \frac{2}{9}$, which implies $m_{Ind_s}(\ell) = 1$ and $\alpha(s) = 1$, then ε is $g_s^{(\ell)}$ -valid if and only if $-\bar{a}(\Delta_3^*)\varepsilon \leq \frac{2}{9}$, i.e., any $\varepsilon > 0$ is $g_s^{(\ell)}$ -valid.

- If $g_s^{(\ell)}(\mathcal{C}_3^*) = \frac{3}{9}$, which implies $m_{Ind_s}(\ell) = 2$ and $\alpha(s) = 3$, then ε is $g_s^{(\ell)}$ -valid if and only if

$$\begin{cases} -(2\bar{a}(\Delta_3^*) + \bar{y}(\Delta_3^*))\varepsilon \leq \frac{3}{9}, \\ -(2\bar{a}(\Delta_3^*) + \bar{b}(\Delta_3^*))\varepsilon \leq \frac{3}{9}, \\ -(\bar{x}(\Delta_3^*) + 2\bar{y}(\Delta_3^*))\varepsilon \leq \frac{3}{9}, \\ -(\bar{x}(\Delta_3^*) + 2\bar{b}(\Delta_3^*))\varepsilon \leq \frac{3}{9}, \\ -(\bar{x}(\Delta_3^*) + \bar{b}(\Delta_3^*) + \bar{y}(\Delta_3^*))\varepsilon \leq \frac{3}{9}. \end{cases}$$

It then follows that, for this case, $\varepsilon > 0$ is $g_s^{(\ell)}$ -valid if and only if $\varepsilon \leq \frac{3}{9 \max\{3, 12\}} = \frac{1}{36}$.

- If $g_s^{(\ell)}(\mathcal{C}_3^*) = \frac{i}{9}$, where $i \geq 4$, then, similarly as above, ε is $g_s^{(\ell)}$ -valid if and only if $\varepsilon > 0$ satisfies the following systems of inequalities:

$$\begin{cases} d_1\varepsilon \leq \frac{i}{9}, \\ d_2\varepsilon \leq \frac{i}{9}, \\ \dots \\ d_r\varepsilon \leq \frac{i}{9}, \end{cases}$$

for some integer r . It is easy to see that for all $1 \leq j \leq r$, $d_j \leq -(\bar{x}(\Delta_3^*) + 2\bar{y}(\Delta_3^*)) = 12$. So, for this case, we have $\varepsilon \leq \frac{1}{36}$ is $g_s^{(\ell)}$ -valid.

Combining all the discussions as above, we conclude that $\varepsilon > 0$ is $g_{\mathcal{S}_k}$ -valid if and only if $\varepsilon \leq \frac{1}{36}$, which completes the proof.

B Proof of Lemma 5.9

Firstly, note that for any 4-sample s and any $1 \leq \ell \leq 4$,

$$g_s^{(\ell)}(\mathcal{C}_4^*) = \frac{4}{16}m_{Ind_s}(\ell) - \frac{1}{16}\alpha(s), \quad (55)$$

whence we have

$$\{g_s^{(\ell)}(\mathcal{C}_4^*) \mid s \in \mathcal{S}_4, 1 \leq \ell \leq 4\} \subseteq \left\{ \frac{-3}{16}, \frac{-2}{16}, \dots, \frac{15}{16}, 1 \right\}.$$

We now consider the following cases:

- If $g_s^{(\ell)}(\mathcal{C}_4^*) = \frac{-3}{16}$, which implies $m_{Ind_s}(\ell) = 0$ and $\alpha(s) = 3$ (see Equation (55)), then ε is $g_s^{(\ell)}$ -valid if and only if

$$\begin{cases} 3\bar{y}(\Delta_4^*)\varepsilon \leq \frac{3}{16}, \\ 3\bar{b}(\Delta_4^*)\varepsilon \leq \frac{3}{16}, \\ (2\bar{y}(\Delta_4^*) + \bar{b}(\Delta_4^*))\varepsilon \leq \frac{3}{16}, \\ (\bar{y}(\Delta_4^*) + 2\bar{b}(\Delta_4^*))\varepsilon \leq \frac{3}{16}. \end{cases}$$

Noting from Example 5.5 that $\bar{x}(\Delta_3) = 3$, $\bar{y}(\Delta_4^*) = -5$, $\bar{a}(\Delta_4^*) = -1$ and $\bar{b}(\Delta_4^*) = 3$, we have that, for this case, $\varepsilon > 0$ is $g_s^{(\ell)}$ -valid if and only if $3\bar{b}(\Delta_4^*)\varepsilon \leq \frac{3}{16}$, i.e., $\varepsilon \leq \frac{1}{48}$.

- If $g_s^{(\ell)}(\mathcal{C}_4^*) = \frac{-2}{16}$, which implies $m_{Ind_s}(\ell) = 0$ and $\alpha(s) = 2$, then ε is $g_s^{(\ell)}$ -valid if and only if

$$\begin{cases} 2\bar{y}(\Delta_4^*)\varepsilon \leq \frac{2}{16}, \\ 2\bar{b}(\Delta_4^*)\varepsilon \leq \frac{2}{16}, \\ \bar{y}(\Delta_4^*)\varepsilon + \bar{b}(\Delta_4^*)\varepsilon \leq \frac{2}{16}. \end{cases}$$

Similarly as above, we deduce that, for this case, $\varepsilon > 0$ is $g_s^{(\ell)}$ -valid if and only if $2\bar{b}(\Delta_4^*)\varepsilon \leq \frac{2}{16}$, i.e., $\varepsilon \leq \frac{1}{48}$.

- If $g_s^{(\ell)}(\mathcal{C}_4^*) = \frac{-1}{16}$, which implies $m_{Ind_s}(\ell) = 0$ and $\alpha(s) = 1$, then ε is $g_s^{(\ell)}$ -valid if and only if $\bar{b}(\Delta_4^*)\varepsilon \leq \frac{1}{16}$, i.e., $\varepsilon \leq \frac{1}{48}$.
- If $g_s^{(\ell)}(\mathcal{C}_4^*) = 0$, then by definition, any $\varepsilon > 0$ is $g_s^{(\ell)}$ -valid.
- If $g_s^{(\ell)}(\mathcal{C}_4^*) = \frac{1}{16}$, which implies $m_{Ind_s}(\ell) = 1$ and $\alpha(s) = 3$, then ε is $g_s^{(\ell)}$ -valid if and only if

$$\begin{cases} -(\bar{a}(\Delta_4^*) + 2\bar{y}(\Delta_4^*))\varepsilon \leq \frac{1}{16}, \\ -(\bar{a}(\Delta_4^*) + 2\bar{b}(\Delta_4^*))\varepsilon \leq \frac{1}{16}, \\ -(\bar{a}(\Delta_4^*) + \bar{b}(\Delta_4^*) + \bar{y}(\Delta_4^*))\varepsilon \leq \frac{1}{16}. \end{cases}$$

We then deduce that, for this case, $\varepsilon > 0$ is $g_s^{(\ell)}$ -valid if and only if $-(\bar{a}(\Delta_4^*) + 2\bar{y}(\Delta_4^*))\varepsilon \leq \frac{1}{16}$, i.e., $\varepsilon \leq \frac{1}{176}$.

- If $g_s^{(\ell)}(\mathcal{C}_4^*) = \frac{2}{16}$, which implies $m_{Ind_s}(\ell) = 1$ and $\alpha(s) = 2$, then ε is $g_s^{(\ell)}$ -valid if and only if

$$\begin{cases} -(\bar{a}(\Delta_4^*) + \bar{y}(\Delta_4^*))\varepsilon \leq \frac{2}{16}, \\ -(\bar{a}(\Delta_4^*) + \bar{b}(\Delta_4^*))\varepsilon \leq \frac{2}{16}. \end{cases}$$

We then infer that $\varepsilon > 0$ is $g_s^{(\ell)}$ -valid if and only if $-(\bar{a}(\Delta_4^*) + \bar{y}(\Delta_4^*))\varepsilon \leq \frac{2}{16}$, i.e., $\varepsilon \leq \frac{1}{48}$.

- If $g_s^{(\ell)}(\mathcal{C}_4^*) = \frac{3}{16}$, which implies $m_{Ind_s}(\ell) = 1$ and $\alpha(s) = 1$, then ε is $g_s^{(\ell)}$ -valid if and only if $-\bar{a}(\Delta_4^*)\varepsilon \leq \frac{3}{16}$, i.e., any $\varepsilon > 0$ is valid.

- If $g_s^{(\ell)}(\mathcal{C}_4^*) = \frac{i}{16}$, where $i = 4, 5, \dots, 16$, then similarly as above, ε is $g_s^{(\ell)}$ -valid if and only if $\varepsilon > 0$ satisfies the following system of inequalities,

$$\begin{cases} d_1\varepsilon \leq \frac{i}{16}, \\ d_2\varepsilon \leq \frac{i}{16}, \\ \dots \\ d_r\varepsilon \leq \frac{i}{16}, \end{cases}$$

for some integer r . It is easy to see that for all $1 \leq j \leq r$, $d_j \leq -(\bar{a}(\mathbf{\Delta}_4^*) + 3\bar{y}(\mathbf{\Delta}_4^*)) = 16$. It then follows that, for this case, $\varepsilon \leq \frac{1}{64}$ is $g_s^{(\ell)}$ -valid.

Combining all the discussions above, we conclude that $\varepsilon > 0$ is g_{S_k} -valid if and only if $\varepsilon \leq \frac{1}{176}$, which completes the proof.

C Proof of Lemma 6.8

For any $\mathcal{C} \in N(\mathcal{C}_4^{**}, \varepsilon)$, we write

$$\mathbf{\Delta} = \mathcal{C} - \mathcal{C}_4^{**} = \left(\left(\delta_{i,j}^{(1)} \right), \left(\delta_{i,j}^{(2)} \right), \left(\delta_{i,j}^{(3)} \right), \left(\delta_{i,j}^{(4)} \right) \right), \quad h_s(\mathbf{\Delta}) = g_s(\mathcal{C}) - g_s(\mathcal{C}_4^{**}).$$

By Lemmas 6.1 and 6.2, we have

$$\sum_{s \in \mathcal{S}_4(3,3,0)} h_s(\mathbf{\Delta}) = -16 \sum_{i=1}^4 \delta_{i,i}^{(i)}, \quad \sum_{s \in \mathcal{S}_4(3,3,3)} h_s(\mathbf{\Delta}) = 3 \sum_{i=1}^4 \delta_{i,i}^{(i)} + \sum_{i \neq j} \delta_{i,i}^{(j)}.$$

We then have the following cases:

- For the samples in $\mathcal{S}_4(3,3,1)$, we write

$$\sum_{s \in \mathcal{S}_4(3,3,1)} h_s(\mathbf{\Delta}) = \sum_{\ell=1}^4 \sum_{i=1}^4 \sum_{j=1}^4 k_{i,j}^{(\ell)} \delta_{i,j}^{(\ell)}, \quad (56)$$

where the coefficients $k_{i,j}^{(\ell)}$ can be determined as follows. First, we consider $(i, i) \in [4] \times [4]$; and for simplicity only, assume $i = 1$. There are 12 samples from $\mathcal{S}_4(3,3,1)$ containing $(1, 1)$, more precisely, those samples from

$$\begin{aligned} & \{ \{(1, 1), (i, 2), (j, 3)\} \mid i = 1, 3; j = 1, 2\} \cup \{ \{(1, 1), (i, 2), (j, 4)\} \mid i = 1, 4; j = 1, 2\} \\ & \cup \{ \{(1, 1), (i, 3), (j, 4)\} \mid i = 1, 4; j = 1, 3\}. \end{aligned}$$

By (14), it is easy to see that $k_{1,1}^{(1)} = 12$ and $k_{1,1}^{(2)} = k_{1,1}^{(3)} = k_{1,1}^{(4)} = 4$. The other coefficients of $\delta_{i,i}^{(\ell)}$ can be obtained similarly as

- $k_{i,i}^{(i)} = 12$, $1 \leq i \leq 4$;
- $k_{i,i}^{(j)} = 4$, $i \neq j$.

We now consider $(i, j) \in [4] \times [4]$ for $i \neq j$; and for simplicity only, assume $(i, j) = (2, 1)$. There are 8 samples of $\mathcal{S}_4(3, 3, 1)$ containing $(2, 1)$, more precisely, those samples from

$$\begin{aligned} & \{ \{(2, 1), (2, 2), (i, 3)\} \mid i = 1, 2\} \cup \{ \{(2, 1), (i, 2), (3, 3)\} \mid i = 1, 3\} \\ & \cup \{ \{(2, 1), (2, 2), (i, 4)\} \mid i = 1, 2\} \cup \{ \{(2, 1), (i, 2), (4, 4)\} \mid i = 1, 4\}. \end{aligned}$$

By (14), it is easy to see that $k_{2,1}^{(1)} = k_{2,1}^{(2)} = 8$ and $k_{2,1}^{(3)} = k_{2,1}^{(4)} = 0$. The other coefficients of $\delta_{i,j}^{(\ell)}$ can be obtained similarly as

- $k_{i,j}^{(i)} = k_{j,i}^{(i)} = 8, 1 \leq i, j \leq 4$;
- $k_{i,j}^{(\ell)} = 0$, if i, j, ℓ are distinct.

Finally, by (36), we have

$$\sum_{s \in \mathcal{S}_4(3,3,1)} h_s(\Delta) = -4 \sum_{i=1}^4 \delta_{i,i}^{(i)} + 4 \sum_{i \neq j} \delta_{i,i}^{(j)}.$$

► For the 81 samples in $\mathcal{S}_4(4, 4, 0)$, as in the previous case, we write

$$\sum_{s \in \mathcal{S}_4(4,4,0)} h_s(\Delta) = \sum_{\ell=1}^4 \sum_{i=1}^4 \sum_{j=1}^4 k_{i,j}^{(\ell)} \delta_{i,j}^{(\ell)}. \quad (57)$$

Note that for $(i, i) \in [4] \times [4]$, since there is no sample containing (i, i) , $k_{i,i}^{(\ell)} = 0$. We then consider $(i, j) \in [4] \times [4]$ for $i \neq j$; and for simplicity only, assume $(i, j) = (2, 1)$. There are 27 samples of $\mathcal{S}_4(4, 4, 0)$ containing $(2, 1)$, more precisely, those samples from

$$\{ \{(2, 1), (i, 2), (j, 3), (\ell, 4)\} \mid i \neq 2, j \neq 3, \ell \neq 4; 1 \leq i, j, \ell \leq 4\}.$$

It is easy to verify that $k_{2,1}^{(1)} = k_{2,1}^{(2)} = k_{2,1}^{(3)} = k_{2,1}^{(4)} = 27$, and the other coefficients of $\delta_{i,j}^{(\ell)}$ can be obtained similarly. All in all, we have $k_{i,j}^{(\ell)} = 27$, for $i \neq j$. Hence, we have

$$\sum_{s \in \mathcal{S}_4(4,4,0)} h_s(\Delta) = -27 \sum_{i=1}^4 \sum_{j=1}^4 \delta_{j,j}^{(i)}.$$

Combining the above results, we have

$$\frac{1}{8} \sum_{s \in \mathcal{S}_3(3,3,0)} h_s(\Delta) + \frac{1}{4} \sum_{s \in \mathcal{S}_4(3,3,1)} h_s(\Delta) + 2 \sum_{s \in \mathcal{S}_4(3,3,3)} h_s(\Delta) + \frac{1}{9} \sum_{s \in \mathcal{S}_4(4,4,0)} h_s(\Delta) = 0, \quad (58)$$

which completes the proof.

References

- [1] R. Ahlswede, N. Cai, S. -Y. R. Li and R. Yeung, “Network information flow,” *IEEE Trans. Inf. Theory*, vol. 46, no. 4, pp. 1204-1216, Aug. 2000.
- [2] C. K. Ngai and R. W. Yeung, “Network coding gain of combination networks,” in *Proc. ITW*, 2004.
- [3] S. Jaggi, P. Sanders, P. A. Chou, M. Effros, S. Egner, K. Jain and L. M. G. Tolhuizen, “Polynomial time algorithms for multicast network code construction,” *IEEE Trans. Inf. Theory*, vol. 51, no. 6, pp. 1973-1982, Jun. 2005.
- [4] A. Agarwal and M. Charikar, “On the advantage of network coding for improving network throughput,” in *Proc. ITW*, 2004.
- [5] Z. Li and B. Li, “Network coding in undirected networks,” in *Proc. 38th CISS*, 2004.
- [6] Z. Li, B. Li and L. C. Lau, “A constant bound on throughput improvement of multicast network coding in undirected networks,” *IEEE Trans. Inf. Theory*, vol. 55, no. 3, pp. 997-1015, Mar. 2009.
- [7] S. Maheshwar, Z. Li and B. Li, “Bounding the coding advantage of combination network coding in undirected networks,” *IEEE Trans. Inf. Theory*, vol. 58, no. 2, pp. 570-584, Feb. 2012.
- [8] Z. Li and B. Li, “Network coding: The case of multiple unicast sessions,” in *Proc. 42nd Annual Allerton*, 2004.
- [9] N. J. A. Harvey, R. D. Kleinberg and A. R. Lehman, “Comparing network coding with multicommodity flow for the k-pairs communication problems,” *Tech. Rep. Comput. Sci. and Artific. Intell. Lab., MIT*, 2004.
- [10] N. Harvey, R. Kleinberg and A. Lehman, “On the capacity of information networks,” *IEEE Trans. Inf. Theory*, vol. 52, no. 6, pp. 2345-2364, Jun. 2006.
- [11] K. Jain, V. V. Vazirani and G. Yuval, “On the capacity of multiple unicast sessions in undirected graphs,” *IEEE/ACM Trans. Networking*, vol. 14, pp. 2805-2809, Jun. 2006.
- [12] T. Xiahou, C. Wu, J. Huang and Z. Li, “A geometric framework for investigating the multiple unicast network coding conjecture,” in *Proc. NetCod*, 2012.
- [13] Z. Li and C. Wu, “Space information flow: multiple unicast,” in *Proc. ISIT 2012*, July 1-6, 2012.
- [14] Y. Yang, X. Yin, X. Chen, Y. Yang and Z. Li, “A note on the multiple-unicast network coding conjecture,” *IEEE Communications Letters*, vol. 18, no. 5, pp. 869-872, May 2014.
- [15] M. Langberg and M. Médard, “On the multiple unicast network coding conjecture,” in *Proc. 47th Annual Allerton*, 2009.
- [16] A. Schrijver, “Combinatorial Optimization,” Springer-Verlag, 2003.

- [17] K. Cai and G. Han, “On the solvability of three-pair networks with common bottleneck links,” in *Proc. ITW*, 2014.
- [18] K. Cai and G. Han, “On network coding advantage for multiple unicast networks,” in *Proc. ISIT*, 2015.
- [19] K. Cai and G. Han, “Coding advantage in communications among peers,” in *Proc. ISIT*, 2016.
- [20] K. Cai and G. Han, “On the Langberg-Médard multiple unicast conjecture,” *Journal of Combinatorial Optimization*, vol. 34, no. 4, pp. 1114-1132, 2017.
- [21] K. Cai and G. Han, “On the Langberg-Médard k -unicast conjecture with $k = 3, 4$,” in *Proc. ISIT*, 2018.